

These are findings from an Ipsos poll conducted for Thomson Reuters April 23-27, 2015. For the survey, a sample of 1,752 adults, including 751 Democrats, 567 Republicans, and 248 Independents ages 18+ were interviewed online. The precision of the Reuters/Ipsos online polls is measured using a [credibility interval](#). In this case, the poll has a credibility interval of plus or minus 2.7 percentage points for all adults, 4.1 for Democrats, 4.7 for Republicans, and 7.1 for Independents. For more information about credibility intervals, please see the appendix.

The data were weighted to the U.S. current population data by gender, age, education, and ethnicity. Statistical margins of error are not applicable to online polls. All sample surveys and polls may be subject to other sources of error, including, but not limited to coverage error and measurement error. Figures marked by an asterisk (*) indicate a percentage value of greater than zero but less than one half of one per cent. Where figures do not sum to 100, this is due to the effects of rounding. To see more information on this and other Reuters/Ipsos polls, please visit <http://polling.reuters.com/>.

SAME-SEX WEDDINGS

Q1. If you were invited to attend a gay wedding of someone you knew, would you attend?

	<u>Total</u>	<u>Democrats</u>	<u>Republicans</u>	<u>Independents</u>
Yes	64%	77%	52%	62%
No	21%	12%	32%	22%
Not sure	15%	10%	16%	16%

Q2. If you were invited to attend the gay wedding for a relative or friend, would you attend?

	<u>Total</u>	<u>Democrats</u>	<u>Republicans</u>	<u>Independents</u>
Yes	66%	80%	53%	67%
No	20%	12%	29%	21%
Not sure	14%	8%	18%	12%

Q3. If you were invited to attend the gay wedding for a loved one, would you attend?

	<u>Total</u>	<u>Democrats</u>	<u>Republicans</u>	<u>Independents</u>
Yes	68%	80%	56%	70%
No	19%	12%	27%	17%
Not sure	13%	8%	16%	12%

How to Calculate Bayesian Credibility Intervals

The calculation of credibility intervals assumes that Y has a binomial distribution conditioned on the parameter θ , i.e., $Y|\theta \sim \text{Bin}(n, \theta)$, where n is the size of our sample. In this setting, Y counts the number of “yes”, or “1”, observed in the sample, so that the sample mean (\bar{y}) is a natural estimate of the true population proportion θ . This model is often called the likelihood function, and it is a standard concept in both the Bayesian and the Classical framework. The Bayesian ¹ statistics combines both the prior distribution and the likelihood function to create a posterior distribution. The posterior distribution represents our opinion about which are the plausible values for θ adjusted after observing the sample data. In reality, the posterior distribution is one’s knowledge base updated using the latest survey information. For the prior and likelihood functions specified here, the posterior distribution is also a beta distribution ($\pi(\theta/y) \sim \beta(y+a, n-y+b)$), but with updated hyper-parameters.

Our credibility interval for ϑ is based on this posterior distribution. As mentioned above, these intervals represent our belief about which are the most plausible values for ϑ given our updated knowledge base. There are different ways to calculate these intervals based on $\pi(\theta/y)$. Since we want only one measure of precision for all variables in the survey, analogous to what is done within the Classical framework, we will compute the largest possible credibility interval for any observed sample. The worst case occurs when we assume that $a=1$ and $b=1$ and $y=n/2$. Using a simple approximation of the posterior by the normal distribution, the 95% credibility interval is given by, approximately:

$$\bar{y} \pm \frac{1}{\sqrt{n}}$$

For this poll, the Bayesian Credibility Interval was adjusted using standard weighting design effect $1+L=1.3$ to account for complex weighting²

Examples of credibility intervals for different base sizes are below. Ipsos does not publish data for base sizes (sample sizes) below 100.

Sample size	Credibility intervals
2,000	2.5
1,500	2.9
1,000	3.5
750	4.1
500	5.0
350	6.0
200	7.9
100	11.2

¹ *Bayesian Data Analysis, Second Edition, Andrew Gelman, John B. Carlin, Hal S. Stern, Donald B. Rubin, Chapman & Hall/CRC | ISBN: 158488388X | 2003*

² Kish, L. (1992). *Weighting for unequal Pi*. *Journal of Official, Statistics*, 8, 2, 183200.