

canprov.....	1
Education Of Panel Member	2
Employment Of Panel Member	3
Household Income.....	4
S1. Please record your gender.....	5
S2. Into which of the following age groups do you belong?.....	6
S3. Do you currently have any children under the age of three living in your household?.....	7
S4. Are you the primary caregiver for a child under three years old in your household?.....	8
Q1. How many children under the age of 18 currently live in your household?	9
Q2. Please record the ages of each of your children, starting with the age of your youngest child.....	10
Q2. Age of the Youngest Child - In MONTHS.....	12
Q3. [SUMMARY - YES] Here are some diseases, please indicate whether you have heard of a vaccine for each of them.	13
Q4. [SUMMARY - YES] Do you think the vaccines for the following diseases are available for children under the age of 2?	14
Q5. [SUMMARY - YES] Please indicate whether your youngest child has received vaccination ... for each of the following diseases.....	15
Q6. [SUMMARY - TOP2BOX] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?.....	16
Q6. [SUMMARY - LOW2BOX] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?.....	17
Q6_1. [Runny nose] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?	18
Q6_2. [Fever] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?	19
Q6_3. [Cough] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?	20
Q6_4. [Mild diarrhea] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?	21
Q6_5. [Forceful vomiting with increased frequency] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?	22
Q6_6. [Decreased activity] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?	23
Q6_7. [Hives] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?	24
Q6_8. [Severe diarrhea] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?	25
Q7. Have you ever heard of a virus called rotavirus, which causes gastroenteritis ... ?	26
Q8. How familiar would you say you are with the rotavirus —very familiar, familiar, not too familiar, or not at all familiar?.....	27
Q9. Has your youngest child received vaccination ... for the rotavirus?	28
Q10. [SUMMARY - TOP2BOX] Here is a list of conditions that could affect your child. Thinking about your youngest child, please indicate how serious you think each condition is—very serious, serious, not too serious or not at all serious	29
Q10. [SUMMARY - LOW2BOX] Here is a list of conditions that could affect your child. Thinking about your youngest child, please indicate how serious you think each condition is—very serious, serious, not too serious or not at all serious	30
Q10_1. [Rotavirus] Here is a list of conditions that could affect your child. Thinking about your youngest child, please indicate how serious you think each condition is—very serious, serious, not too serious or not at all serious.	31
Q10_2. [Influenza] Here is a list of conditions that could affect your child. Thinking about your youngest child, please indicate how serious you think each condition is—very serious, serious, not too serious or not at all serious.....	32
Q10_3. [Whooping Cough] Here is a list of conditions that could affect your child. Thinking about your youngest child, please indicate how serious you think each condition is—very serious, serious, not too serious or not at all serious.	33

Q10_4. [Chicken Pox] Here is a list of conditions that could affect your child. Thinking about your youngest child, please indicate how serious you think each condition is—very serious, serious, not too serious or not at all serious.	34
Q10_5. [Measles] Here is a list of conditions that could affect your child. Thinking about your youngest child, please indicate how serious you think each condition is—very serious, serious, not too serious or not at all serious.	35
Q11. [SUMMARY - YES] Here are the same conditions. For each, please indicate if you have seen, read, or heard anything about it in the past 6 months.	36
Q12. From what sources have you heard about the rotavirus ... ?	37
Q13. Please describe the information you saw or heard about the rotavirus ... ?	39
Q14. Has your youngest child ever experienced five or more episodes of profuse, watery diarrhea in one day?	41
Q15. How many times has your youngest child experienced this in his/ her lifetime?	42
Q16. Has/ have your other child/ children ever experienced five or more episodes of profuse, watery diarrhea in one day?	43
Q17. Thinking about the last time your youngest child had five or more episodes of profuse, watery diarrhea in one day, did you ...?	44
Q18. Did your child stay overnight at a hospital ... ?	45
Q19. Has your youngest child or any of your children ever stayed for at least one night at a hospital because of multiple episodes of profuse and watery diarrhea?	46
Q20. How well do you feel you managed the last incidence of profuse and watery diarrhea with your youngest child?	47
Q21. The first time your youngest child experienced multiple episodes of profuse, watery diarrhea, how worried were you—very worried, worried, not too worried or not at all worried?	48
Q22. How serious do you find this condition to be?	49
Q23. To the best of your knowledge, which, if any, of the following are signs of dehydration among infants?	50
Q24. Has your youngest child ever been diagnosed by a doctor with viral diarrhea or rotavirus?	51
Q25. Has/ have your other child/ children ever been diagnosed by a doctor with viral diarrhea or rotavirus?	52
Q26. Which, if any, of the following are possible symptoms or consequences of viral diarrhea or rotavirus?	53
Q27. [SUMMARY - TRUE] Now, please indicate if you think the following statements about rotavirus or viral diarrhea are true or false.	54
Q27. [SUMMARY - FALSE] Now, please indicate if you think the following statements about rotavirus or viral diarrhea are true or false.	55
Q28. Besides your child's primary healthcare provider, from which two of the following sources do you obtain most of your information about health issues for your children?	56
Q29. Based on your two previous responses, can you name specific sources where you obtain your information about health issues for your children?	58
Q30. How would you describe the area where you live -- is it located in a city, in a suburb outside of a city, in a small town, or in a country or farm area?	61
Q14/16. Has your child ever experienced five or more episodes of profuse, watery diarrhea in one day?	62
Q24/Q25. Has your child ever been diagnosed by a doctor with viral diarrhea or rotavirus?	63

Rotavirus Study (60-6056-01)

canprov.

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All Respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
British Columbia	110	110	0	0	0	0	39	71	63	47	9	101	36	74	45	65
	13%	100% BCDE	-	-	-	-	15%	13%	16% I	11%	32%	13%	15%	13%	15%	12%
Alberta	84	0	84	0	0	0	28	56	41	43	3	81	22	63	25	60
	10%	-	59% ACDE	-	-	-	11%	10%	10%	10%	11%	10%	9%	11%	8%	11%
Manitoba/Saskatchewan	58	0	58	0	0	0	16	42	23	36	1	57	13	45	22	37
	7%	-	41% ACDE	-	-	-	6%	8%	6%	8%	4%	7%	6%	8%	7%	7%
Ontario	303	0	0	303	0	0	88	215	137	166	9	294	78	225	100	203
	37%	-	-	100% ABDE	-	-	34%	38%	35%	39%	31%	37%	32%	39%	33%	39%
Quebec	200	0	0	0	200	0	69	130	93	106	5	195	74	125	80	119
	24%	-	-	-	100% ABCE	-	26%	23%	24%	25%	17%	25%	31% M	22%	27%	23%
Atlantic	67	0	0	0	0	67	22	45	37	31	2	65	17	50	27	40
	8%	-	-	-	-	100% ABCD	8%	8%	9%	7%	6%	8%	7%	9%	9%	8%

Rotavirus Study (60-6056-01)

Education Of Panel Member

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All Respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Primary School or less	3 0	0 -	0 -	1 0	2 1%	0 -	1 0	2 0	0 -	3 1%	0 -	3 0	0 -	3 1%	1 0	2 0
Some High School	92 11%	5 5%	17 12%	31 10%	36 18%	2 3%	19 7%	73 13%	34 9%	58 13%	2 7%	90 11%	22 9%	69 12%	40 13%	52 10%
High School	244 30%	37 34%	46 32%	94 31%	48 24%	19 29%	79 30%	165 29%	99 25%	145 34%	8 28%	236 30%	62 26%	182 31%	93 31%	152 29%
Some Community College/CEGEP/Trade School	276 34%	48 44%	53 37%	94 31%	59 30%	21 31%	83 32%	192 34%	129 33%	147 34%	13 45%	262 33%	72 30%	204 35%	99 33%	176 34%
Community College/CEGEP/Trade School	27 3%	3 3%	2 1%	9 3%	5 3%	8 12%	10 4%	18 3%	14 4%	13 3%	2 7%	25 3%	8 3%	19 3%	12 4%	15 3%
Some University	111 13%	12 11%	16 12%	39 13%	32 16%	10 16%	45 17%	66 12%	70 18%	41 10%	3 10%	108 14%	45 19%	66 11%	36 12%	75 14%
University Undergraduate degree	56 7%	2 2%	6 4%	30 10%	11 6%	7 10%	21 8%	36 6%	38 10%	18 4%	0 -	56 7%	25 10%	32 5%	15 5%	42 8%
University Graduate degree	13 2%	2 2%	1 1%	4 1%	6 3%	0 -	4 2%	9 2%	9 2%	4 1%	1 3%	12 2%	7 3%	6 1%	3 1%	10 2%

Rotavirus Study (60-6056-01)

Employment Of Panel Member

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All Respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Employed Full-time	224 27%	21 19%	25 17%	83 27% B	77 39% ABC	19 29%	85 32% G	140 25%	113 29%	112 26%	7 24%	217 27%	69 29%	155 27%	68 23%	156 30% N
Employed Part-time	83 10%	13 12% D	16 12% D	34 11% D	9 5%	10 14% D	27 10%	56 10%	28 7%	55 13% H	2 7%	81 10%	14 6%	69 12% L	33 11%	50 10%
Self-Employed	45 6%	10 9% C	12 9% C	9 3%	12 6%	2 3%	12 5%	33 6%	28 7% I	17 4%	4 14%	41 5%	17 7%	28 5%	27 9% O	18 3%
Homemaker	362 44%	57 52% D	72 50% D	141 47% D	66 33%	25 38%	97 37%	266 47% F	174 44%	188 44%	13 45%	349 44%	107 44%	255 44%	133 45%	229 44%
Student	19 2%	1 1%	7 5% C	4 1%	7 4%	0 -	5 2%	14 3%	10 3%	9 2%	0 -	19 2%	4 2%	15 3%	8 3%	11 2%
Retired	1 0	1 1%	0 -	0 -	0 -	0 -	1 0	0 -	1 0	0 -	1 4%	0 -	0 -	1 0	0 -	1 0
Currently Unemployed	34 4%	4 4%	2 1%	17 6% B	7 4%	3 5%	15 6%	19 3%	11 3%	23 5%	2 7%	32 4%	8 3%	26 4%	15 5%	19 4%
Other	52 6%	2 2%	8 6%	13 4%	21 11% AC	8 12% AC	21 8%	32 6%	27 7%	25 6%	0 -	52 7%	20 8%	32 5%	13 4%	39 8%
(Dk/Ns)	1 0	0 -	0 -	1 0	0 -	0 -	0 -	1 0	1 0	0 -	0 -	1 0	1 0	0 -	1 0	0 -

Rotavirus Study (60-6056-01)

Household Income

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All Respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
<\$20K	81 10%	11 10%	13 9%	25 8%	24 12%	7 10%	31 12%	50 9%	32 8%	49 11%	1 4%	80 10%	19 8%	62 11%	31 10%	50 10%
\$20,000-\$29,999	93 11%	11 10%	14 10%	36 12%	21 11%	10 14%	27 10%	66 12%	37 9%	56 13%	6 20%	87 11%	23 10%	70 12%	37 12%	56 11%
\$30,000-\$39,999	87 11%	17 16% C	17 12% C	16 5%	24 12% C	12 18% C	33 13%	54 10%	36 9%	52 12%	3 10%	84 11%	24 10%	63 11%	37 12%	50 10%
\$40,000-\$54,999	125 15%	24 21% C	26 18% C	27 9%	38 19% C	10 16% C	34 13%	91 16%	57 15%	68 16%	4 14%	121 15%	33 14%	92 16%	40 13%	85 16%
\$55,000-\$59,999	53 6%	4 4%	12 9%	18 6%	14 7%	4 6%	15 6%	38 7%	22 5%	31 7%	0 -	53 7%	18 7%	35 6%	22 7%	31 6%
\$60,000-\$69,999	105 13%	16 15%	15 11%	41 14%	26 13%	6 9%	30 11%	75 13%	60 15% I	45 11%	6 20%	99 13%	30 12%	75 13%	36 12%	69 13%
\$70,000-\$79,999	63 8%	6 6%	15 11%	22 7%	12 6%	7 10%	23 9%	40 7%	37 9%	26 6%	2 7%	61 8%	29 12% M	34 6%	25 8%	38 7%
\$80,000-\$99,999	77 9%	6 6%	9 6%	45 15% ABD	11 6%	5 8%	22 8%	55 10%	34 9%	43 10%	4 14%	73 9%	17 7%	60 10%	25 8%	52 10%
\$100,000+	138 17%	13 12%	19 14%	71 23% ABDE	29 15%	5 8%	47 18%	91 16%	79 20% I	58 14%	3 10%	135 17%	47 20%	91 16%	46 15%	92 18%
Mean	60964. 4	54943. 9	58446. 0	69233. 3 ABDE	56527. 6	52064. 9	60677. 0	61098. 9	65214. 2 I	57063. 7	59130. 3	61031. 9	64551. 7	59483. 4	59022. 9	62071. 7

Rotavirus Study (60-6056-01)

S1. Please record your gender.

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Female	822	110	143	303	200	67	262	560	393	429	29	793	240	582	299	523
	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%

Rotavirus Study (60-6056-01)

S2. Into which of the following age groups do you belong?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Between 20 and 24	82 10%	8 7%	22 15% C	26 9%	21 11%	5 8%	39 15% G	43 8%	31 8%	51 12% H	2 7%	80 10%	22 9%	60 10%	32 11%	50 10%
Between 25 and 29	243 30%	34 31%	46 32%	83 27%	66 33% E	14 21%	97 37% G	145 26%	106 27%	137 32%	12 41%	231 29%	74 31%	169 29%	84 28%	159 30%
Between 30 and 34	286 35%	36 33%	41 29%	102 34%	81 41% B	26 39%	70 27%	216 39% F	158 40% I	128 30%	7 24%	279 35%	92 38%	195 33%	108 36%	178 34%
Between 35 and 39	157 19%	23 21%	22 15%	68 22% D	26 13%	19 29% BD	36 14%	121 22% F	74 19%	83 19%	7 24%	150 19%	38 16%	119 21%	56 19%	101 19%
Between 40 and 45	53 6%	9 8% D	12 9% D	24 8% D	5 3%	3 4%	19 7%	34 6%	24 6%	29 7%	1 4%	52 7%	15 6%	38 7%	18 6%	35 7%
Summary																
Mean (in Years)	31.2	31.6 D	30.5	31.7 BD	30.2	32.0 D	30.1	31.7 F	31.5	30.9	30.8	31.2	31.0	31.2	31.1	31.2

Rotavirus Study (60-6056-01)

S3. Do you currently have any children under the age of three living in your household?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
 Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Yes	822	110	143	303	200	67	262	560	393	429	29	793	240	582	299	523
	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%

Rotavirus Study (60-6056-01)

S4. Are you the primary caregiver for a child under three years old in your household?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Yes	822	110	143	303	200	67	262	560	393	429	29	793	240	582	299	523
	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%

Rotavirus Study (60-6056-01)

Q1. How many children under the age of 18 currently live in your household?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
1	262 32%	39 36%	44 31%	88 29%	69 35%	22 32%	262 100% G	0 -	121 31%	141 33%	15 51%	247 31%	84 35%	178 31%	66 22%	196 37% N
2	347 42%	45 41%	54 38%	134 44%	80 40%	33 49%	0 -	347 62% F	171 44%	176 41%	8 28%	339 43%	97 40%	250 43%	134 45%	213 41%
3	149 18%	16 15%	32 22%	61 20%	32 16%	8 12%	0 -	149 27% F	69 18%	79 19%	0 -	149 19%	42 17%	107 18%	65 22% O	83 16%
4	46 6%	7 7%	9 6%	15 5%	12 6%	3 4%	0 -	46 8% F	23 6%	23 5%	4 14%	42 5%	16 7%	30 5%	25 8% O	21 4%
5	11 1%	1 1%	3 2%	2 1%	4 2%	1 1%	0 -	11 2% F	5 1%	6 1%	1 3%	10 1%	1 0	10 2%	4 1%	7 1%
6	4 0	1 1%	0 -	2 1%	0 -	1 1%	0 -	4 1%	3 1%	1 0	1 4%	3 0	1 0	3 0	2 1%	2 0
7	1 0	0 -	0 -	0 -	1 1%	0 -	0 -	1 0	0 -	1 0	0 -	1 0	0 -	1 0	1 0	0 -
8	2 0	0 -	0 -	1 0	1 1%	0 -	0 -	2 0	1 0	1 0	0 -	2 0	0 -	2 0	1 0	1 0
Summary																
Mean	2.1	2.0	2.1	2.1	2.1	2.0	1.0	2.5 F	2.1	2.0	2.0	2.1	2.0	2.1	2.3 O	1.9
Std Dev	1.0	1.0	1.0	1.0	1.1	1.0	0.0	0.9	1.0	1.0	1.4	1.0	0.9	1.0	1.1	1.0
Std Err	0	0.1	0.1	0.1	0.1	0.1	0.0	0	0.1	0	0.3	0	0.1	0	0.1	0
Median	1.4	1.4	1.5	1.5	1.4	1.4	1.0	2.0	1.4	1.4	1.0	1.4	1.4	1.5	1.6	1.3

Rotavirus Study (60-6056-01)

Q2. Please record the ages of each of your children, starting with the age of your youngest child.

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Have Children 1 - 12 months old	490 60%	49 45%	84 59%	186 61%	129 65%	41 61%	156 60%	334 60%	244 62%	245 57%	19 66%	470 59%	165 68%	325 56%	168 56%	322 61%
Have Children 13 - 24 months old	268 33%	50 46%	50 35%	94 31%	53 27%	20 30%	76 29%	192 34%	125 32%	142 33%	12 41%	256 32%	63 26%	205 35%	110 37%	158 30%
Have Children 25 - 36 months old	83 10%	13 12%	9 6%	33 11%	20 10%	7 10%	28 11%	55 10%	33 8%	50 12%	0 -	83 10%	18 7%	65 11%	34 11%	49 9%
Have Children 37 - 48 months old	21 3%	1 1%	3 2%	9 3%	6 3%	2 3%	1 0	20 4%	9 2%	12 3%	0 -	21 3%	6 3%	15 3%	11 4%	10 2%
Have Children 49 - 60 months old	23 3%	2 2%	4 3%	7 2%	8 4%	2 3%	1 0	22 4%	10 2%	13 3%	1 4%	22 3%	3 1%	20 3%	13 4%	10 2%
Have Children 61 - 72 months old	26 3%	7 7%	4 3%	11 4%	4 2%	0 -	0 -	26 5%	13 3%	13 3%	1 3%	25 3%	9 4%	17 3%	9 3%	17 3%
Have Children 73 - 84 months old	42 5%	5 5%	9 6%	12 4%	10 5%	5 8%	0 -	42 7%	21 5%	21 5%	1 3%	41 5%	10 4%	32 5%	17 6%	25 5%
Have Children 85 - 96 months old	25 3%	3 3%	8 6%	9 3%	2 1%	3 4%	0 -	25 4%	12 3%	13 3%	0 -	25 3%	7 3%	18 3%	13 4%	12 2%
Have Children 97 - 108 months old	45 6%	5 5%	8 6%	17 6%	13 7%	2 3%	9 3%	36 6%	24 6%	21 5%	2 7%	43 5%	15 6%	30 5%	13 4%	32 6%
Have Children 109 - 120 months old	38 5%	3 3%	7 5%	13 4%	13 7%	2 3%	0 -	38 7%	18 5%	20 5%	2 7%	36 5%	11 5%	27 5%	16 5%	22 4%
Have Children 121 - 132 months old	23 3%	2 2%	4 3%	5 2%	10 5%	2 3%	0 -	23 4%	15 4%	8 2%	0 -	23 3%	6 3%	17 3%	11 4%	12 2%
Have Children 133 - 144 months old	12 1%	0 -	4 3%	5 2%	3 2%	0 -	0 -	12 2%	4 1%	8 2%	0 -	12 2%	2 1%	10 2%	5 2%	7 1%
Have Children 145 - 156 months old	16 2%	2 2%	2 1%	6 2%	4 2%	2 3%	0 -	16 3%	7 2%	9 2%	1 4%	15 2%	5 2%	11 2%	6 2%	10 2%
Have Children 157 - 168 months old	9 1%	1 1%	0 -	4 1%	2 1%	2 3%	0 -	9 2%	2 0	7 2%	0 -	9 1%	0 -	9 2%	3 1%	6 1%

Rotavirus Study (60-6056-01)

Q2. Please record the ages of each of your children, starting with the age of your youngest child.

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
Have Children 169 - 180 months old	5	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
	1%	0	1	3	1	0	0	5	0	5	0	5	0	5	3	2
Have Children 181 - 192 months old	10	-	1%	1%	1%	-	-	1%	-	1%	-	1%	-	1%	1%	0
	1%	0	1	5	3	1	0	10	7	3	1	9	4	6	6	4
Have Children 193 - 204 months old	242	27	36	94	69	17	82	161	123	119	11	231	86	156	101	141
	29%	24%	25%	31%	35%	25%	31%	29%	31%	28%	38%	29%	36%	27%	34%	27%
Have Children 205 - 216 months old	75	10	15	28	15	6	0	75	32	43	3	72	13	62	33	42
	9%	9%	11%	9%	8%	9%	-	13%	8%	10%	10%	9%	5%	11%	11%	8%
								F					L			

Rotavirus Study (60-6056-01)

Q2. Age of the Youngest Child - In MONTHS

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted																
	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
IN MONTHS																
1 - 5	91 11%	9 8%	13 9%	34 11%	27 14%	7 10%	22 8%	69 12%	46 12%	45 10%	2 7%	89 11%	33 14%	58 10%	24 8%	67 13% N
6 - 10	138 17%	9 8%	28 19% A	54 18% A	33 17% A	14 21% A	47 18%	91 16%	69 17%	70 16%	4 14%	134 17%	50 21%	89 15%	41 14%	98 19%
11 - 15	259 32%	31 28%	43 30%	97 32%	68 34%	20 30%	87 33%	172 31%	130 33%	129 30%	13 45%	246 31%	82 34%	178 31%	102 34%	157 30%
21 - 25	259 31%	48 44% CDE	48 34%	92 30%	51 26%	19 29%	76 29%	183 33%	121 31%	138 32%	10 34%	249 31%	60 25%	199 34% L	103 34%	156 30%
31+	75 9%	12 11%	10 7%	25 8%	20 10%	7 10%	30 11%	45 8%	28 7%	47 11%	0 -	75 9%	16 7%	59 10%	29 10%	46 9%
Summary																
Mean (in months)	16.4	19.1 BCD	16.3	15.9	15.9	16.3	17.0	16.2	15.6	17.1 H	15.0	16.5	14.5	17.2 L	17.3	15.9
Std Dev	9.9	10.2	9.4	9.5	10.4	9.7	10.1	9.8	9.2	10.4	7.0	10.0	9.1	10.1	9.6	10.0
Std Err	0.3	1.0	0.8	0.6	0.7	1.1	0.6	0.4	0.5	0.5	1.3	0.4	0.6	0.4	0.6	0.4
Median	11.7	13.4 C	11.7	11.6	11.5	11.6 BCD	11.7	11.7	11.6	11.8	11.0	11.7	11.4 M	11.8	11.8	11.6

Rotavirus Study (60-6056-01)

Q3. [SUMMARY - YES] Here are some diseases, please indicate whether you have heard of a vaccine for each of them.

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Measles, mumps and rubella	814 99%	110 100%	141 99%	300 99%	197 98%	67 100%	258 98%	556 99%	391 99%	423 99%	29 100%	785 99%	239 100%	575 99%	294 99%	519 99%
Influenza	800 97%	110 100%	134 94%	293 97%	199 99%	64 96%	252 96%	548 98%	388 99%	412 96%	29 100%	771 97%	237 99%	563 97%	290 97%	510 98%
Diphtheria, whooping cough, tetanus	797 97%	108 98%	139 97%	289 95%	197 98%	65 97%	250 95%	547 98%	387 98%	410 96%	29 100%	768 97%	235 98%	562 97%	287 96%	510 97%
Varicella	787 96%	109 99%	136 96%	290 96%	188 94%	64 96%	246 94%	541 97%	382 97%	405 94%	29 100%	758 96%	232 97%	555 95%	283 95%	504 96%
Hepatitis B	785 96%	108 98%	134 94%	291 96%	190 95%	63 94%	247 94%	538 96%	385 98%	400 93%	29 100%	756 95%	236 98%	549 94%	282 94%	504 96%
Meningitis	781 95%	105 95%	139 97%	289 95%	189 94%	61 91%	248 95%	534 95%	379 96%	402 94%	29 100%	752 95%	232 97%	549 94%	281 94%	501 96%
Polio	770 94%	108 98%	136 96%	295 97%	167 83%	64 96%	238 91%	532 95%	375 95%	395 92%	28 97%	742 94%	228 95%	542 93%	273 92%	497 95%
Pneumonia	665 81%	97 89%	119 83%	256 85%	142 71%	51 75%	213 81%	452 81%	338 86%	327 76%	28 97%	637 80%	204 85%	461 79%	224 75%	441 84%

Rotavirus Study (60-6056-01)

Q4. [SUMMARY - YES] Do you think the vaccines for the following diseases are available for children under the age of 2?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Measles, mumps and rubella	764	109	133	277	188	58	239	525	367	397	28	736	223	541	272	492
	93%	99%	94%	91%	94%	86%	91%	94%	93%	93%	97%	93%	93%	93%	91%	94%
		BCDE			E											
Diphtheria, whooping cough, tetanus	744	105	134	261	183	61	233	511	359	386	27	717	220	524	265	479
	91%	95%	94%	86%	91%	91%	89%	91%	91%	90%	93%	90%	92%	90%	89%	91%
		C	C													
Varicella	686	95	117	251	170	53	221	465	335	351	26	660	213	473	247	440
	84%	87%	82%	83%	85%	79%	84%	83%	85%	82%	90%	83%	89%	81%	83%	84%
													M			
Polio	647	93	107	250	140	56	190	457	316	330	24	623	197	450	232	415
	79%	85%	75%	83%	70%	83%	72%	82%	80%	77%	83%	79%	82%	77%	78%	79%
		D		D		D		F								
Influenza	643	90	122	211	174	46	210	433	326	317	26	617	201	442	228	415
	78%	82%	86%	70%	87%	69%	80%	77%	83%	74%	90%	78%	84%	76%	76%	79%
		CE	CE		CE				I				M			
Meningitis	622	92	106	230	144	51	198	424	309	313	27	595	192	430	229	393
	76%	84%	74%	76%	72%	75%	75%	76%	79%	73%	93%	75%	80%	74%	77%	75%
		D														
Pneumonia	502	79	89	186	111	37	158	344	259	243	23	479	157	345	185	317
	61%	72%	63%	61%	56%	55%	60%	61%	66%	57%	79%	60%	65%	59%	62%	61%
		CDE							I							
Hepatitis B	374	74	48	128	86	37	121	253	185	189	18	356	115	259	139	235
	46%	67%	34%	42%	43%	56%	46%	45%	47%	44%	62%	45%	48%	44%	46%	45%
		BCD				BC										

Rotavirus Study (60-6056-01)

Q5. [SUMMARY - YES] Please indicate whether your youngest child has received vaccination ... for each of the following diseases.

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Diphtheria, whooping cough, tetanus	691 84%	95 87%	121 85%	245 81%	170 85%	60 90%	219 84%	473 84%	333 85%	358 84%	27 93%	664 84%	205 85%	487 84%	258 86%	433 83%
Measles, mumps and rubella	635 77%	100 91% BCDE	106 74%	229 76%	149 74%	52 78%	194 74%	441 79%	291 74%	344 80% H	27 93%	608 77%	173 72%	462 79% L	247 83% O	388 74%
Polio	571 69%	84 77% D	95 67%	212 70%	129 65%	50 74%	167 64%	404 72% F	277 70%	293 68%	19 66%	551 70%	171 71%	400 69%	220 74%	351 67%
Meningitis	475 58%	86 79% BCDE	84 59%	171 56%	97 49%	37 55%	154 59%	321 57%	227 58%	248 58%	24 83%	451 57%	143 60%	332 57%	187 63% O	288 55%
Varicella	422 51%	69 63% CD	78 55% D	154 51% D	82 41% D	38 57% D	139 53%	283 51%	193 49%	228 53%	22 76%	400 50%	126 52%	296 51%	161 54%	261 50%
Pneumonia	397 48%	67 61% CD	82 58% CD	143 47% D	71 36% D	34 51% D	133 51%	264 47%	192 49%	205 48%	20 69%	377 48%	118 49%	279 48%	149 50%	248 47%
Influenza	357 43%	64 58% CD	82 58% CD	99 33% D	74 37% D	38 57% CD	123 47%	234 42%	172 44%	185 43%	22 76%	335 42%	112 46%	246 42%	136 45%	222 42%
Hepatitis B	259 32%	68 62% BCDE	34 24%	88 29% D	42 21% D	28 42% BCD	92 35%	168 30%	128 32%	132 31%	15 52%	244 31%	80 33%	180 31%	93 31%	167 32%

Rotavirus Study (60-6056-01)

Q6. [SUMMARY - TOP2BOX] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Forceful vomiting with increased frequency	799 97%	105 95%	138 96%	297 98%	193 96%	67 100%	254 97%	545 97%	387 98% I	411 96%	28 97%	770 97%	234 97%	565 97%	289 97%	509 97%
Severe diarrhea	791 96%	105 95%	139 97%	294 97%	188 94%	66 99%	253 97%	538 96%	384 98% I	406 95%	29 100%	762 96%	237 99% M	554 95%	285 95%	506 97%
Fever	769 94%	97 89%	132 93%	285 94%	189 94%	66 99% A	248 95%	521 93%	368 94%	401 94%	27 93%	742 94%	228 95%	541 93%	286 96% O	483 92%
Hives	494 60%	76 69% DE	88 62% D	206 68% DE	89 45%	35 52%	159 61%	335 60%	238 60%	256 60%	20 69%	474 60%	149 62%	346 59%	188 63%	307 59%
Decreased activity	448 55%	64 58% D	76 53% D	191 63% D	83 42%	35 52%	138 53%	310 55%	218 55%	231 54%	21 73%	427 54%	141 59%	307 53%	177 59% O	271 52%
Mild diarrhea	385 47%	55 50% B	52 37%	158 52% BD	85 43%	34 51% B	128 49%	257 46%	192 49%	194 45%	17 59%	368 46%	128 53% M	258 44%	130 43%	256 49%
Cough	176 21%	21 19%	28 19%	76 25%	40 20%	12 18%	69 26% G	108 19%	69 18%	107 25% H	7 24%	169 21%	51 21%	125 21%	66 22%	110 21%
Runny nose	44 5%	7 7%	5 4%	22 7%	9 5%	1 1%	20 8% G	24 4%	14 4%	30 7% H	3 10%	41 5%	14 6%	30 5%	14 5%	30 6%

Rotavirus Study (60-6056-01)

Q6. [SUMMARY - LOW2BOX] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

	Total	Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
		BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Runny nose	776 94%	103 93%	136 96%	281 93%	190 95%	66 99%	242 92%	534 95%	378 96%	397 93%	25 86%	750 95%	225 94%	551 95%	284 95%	491 94%
Cough	644 78%	89 81%	114 80%	226 75%	160 80%	55 82%	192 73%	451 81%	323 82%	321 75%	22 76%	622 78%	188 78%	456 78%	232 78%	411 79%
Mild diarrhea	431 52%	54 50%	89 63%	143 47%	111 56%	33 49%	133 51%	299 53%	199 50%	233 54%	11 38%	420 53%	112 46%	320 55%	167 56%	265 51%
Decreased activity	370 45%	46 42%	66 46%	110 36%	115 58%	32 48%	124 47%	246 44%	175 44%	195 45%	7 24%	363 46%	98 41%	272 47%	119 40%	250 48%
Hives	299 36%	33 30%	49 35%	91 30%	96 48%	30 44%	94 36%	205 37%	149 38%	150 35%	8 27%	291 37%	86 36%	213 37%	100 33%	199 38%
Fever	48 6%	12 11%	7 5%	16 5%	11 6%	1 1%	13 5%	35 6%	23 6%	24 6%	2 7%	46 6%	10 4%	37 6%	10 3%	37 7%
Severe diarrhea	27 3%	5 5%	3 2%	9 3%	10 5%	0 -	8 3%	19 3%	7 2%	20 5%	0 -	27 3%	3 1%	24 4%	12 4%	15 3%
Forceful vomiting with increased frequency	20 2%	5 5%	4 3%	5 2%	6 3%	0 -	8 3%	12 2%	5 1%	15 4%	0 -	20 3%	5 2%	15 3%	8 3%	12 2%

Rotavirus Study (60-6056-01)

Q6_1. [Runny nose] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Runny nose																
Very serious	8 1%	0 -	1 1%	7 2% D	0 -	0 -	3 1%	5 1%	5 1%	3 1%	2 7%	6 1%	5 2% M	3 1%	2 1%	6 1%
Serious	36 4%	7 7%	4 3%	15 5%	9 5%	1 1%	17 7% G	19 3%	9 2%	27 6% H	1 4%	35 4%	9 4%	27 5%	12 4%	24 5%
Not too serious	487 59%	65 59%	83 58%	170 56%	121 61%	48 71% C	171 65% G	316 56%	233 59%	253 59%	18 62%	469 59%	144 60%	342 59%	175 59%	312 60%
Not at all serious	289 35%	38 35%	53 37%	111 37%	68 34%	18 27%	71 27%	218 39% F	145 37%	144 34%	7 24%	282 36%	81 34%	208 36%	109 37%	180 34%
(DK/NS)	2 0	0 -	1 1%	0 -	1 1%	0 -	0 -	2 0	1 0	1 0	1 3%	1 0	1 0	1 0	0 -	2 0
Summary																
Top2Box	44 5%	7 7%	5 4%	22 7%	9 5%	1 1%	20 8% G	24 4%	14 4%	30 7% H	3 10%	41 5%	14 6%	30 5%	14 5%	30 6%
Low2Box	776 94%	103 93%	136 96%	281 93%	190 95%	66 99%	242 92%	534 95%	378 96% I	397 93%	25 86%	750 95%	225 94%	551 95%	284 95%	491 94%

Rotavirus Study (60-6056-01)

Q6_2. [Fever] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Fever																
Very serious	506 62%	62 56%	85 60%	196 65%	120 60%	43 64%	184 70% G	322 57%	233 59%	273 64%	20 69%	485 61%	161 67% M	344 59%	182 61%	324 62%
Serious	264 32%	36 33%	47 33%	89 29%	68 34%	24 35%	64 24%	200 36% F	135 34%	128 30%	7 24%	257 32%	67 28%	197 34%	105 35%	159 30%
Not too serious	34 4%	8 7%	5 4%	13 4%	7 4%	1 1%	10 4%	24 4%	20 5%	14 3%	2 7%	32 4%	8 3%	26 5%	8 3%	26 5%
Not at all serious	13 2%	4 4%	2 1%	3 1%	4 2%	0 -	3 1%	10 2%	3 1%	10 2%	0 -	13 2%	2 1%	11 2%	2 1%	11 2%
(DK/NS)	5 1%	0 -	3 2% D	2 1%	0 -	0 -	1 0	4 1%	2 1%	3 1%	0 -	5 1%	2 1%	3 1%	2 1%	3 1%
Summary																
Top2Box	769 94%	97 89%	132 93%	285 94%	189 94%	66 99% A	248 95%	521 93%	368 94%	401 94%	27 93%	742 94%	228 95%	541 93%	286 96% O	483 92%
Low2Box	48 6%	12 11% CE	7 5%	16 5%	11 6%	1 1%	13 5%	35 6%	23 6%	24 6%	2 7%	46 6%	10 4%	37 6%	10 3%	37 7% N

Rotavirus Study (60-6056-01)

Q6_3. [Cough] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Cough																
Very serious	21 3%	1 1%	3 2%	15 5% D	1 1%	1 1%	7 3%	14 3%	9 2%	12 3%	2 7%	19 2%	9 4%	12 2%	8 3%	13 2%
Serious	155 19%	19 18%	25 17%	61 20%	39 20%	11 17%	62 24% G	93 17%	60 15%	95 22% H	5 17%	150 19%	42 18%	113 19%	58 19%	97 19%
Not too serious	541 66%	73 66%	93 65%	188 62%	137 69%	50 74%	159 61%	382 68% F	278 71% I	263 61%	20 69%	521 66%	160 67%	381 66%	193 65%	348 66%
Not at all serious	103 12%	16 15%	21 14%	38 13%	22 11%	5 8%	33 13%	69 12%	45 11%	57 13%	2 7%	101 13%	28 12%	75 13%	39 13%	63 12%
(DK/NS)	2 0	0 -	1 1%	1 0	0 -	0 -	1 0	1 0	1 0	1 0	0 -	2 0	1 0	1 0	0 -	2 0
Summary																
Top2Box	176 21%	21 19%	28 19%	76 25%	40 20%	12 18%	69 26% G	108 19%	69 18%	107 25% H	7 24%	169 21%	51 21%	125 21%	66 22%	110 21%
Low2Box	644 78%	89 81%	114 80%	226 75%	160 80%	55 82%	192 73%	451 81% F	323 82% I	321 75%	22 76%	622 78%	188 78%	456 78%	232 78%	411 79%

Rotavirus Study (60-6056-01)

Q6_4. [Mild diarrhea] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Mild diarrhea																
Very serious	49 6%	2 2%	4 3%	30 10% ABD	6 3%	6 9% ABD	23 9% G	26 5%	22 6%	27 6%	5 17%	43 5%	19 8%	29 5%	15 5%	34 6%
Serious	337 41%	53 49% B	48 34%	128 42%	79 40%	28 42%	105 40%	232 41%	170 43%	167 39%	12 41%	325 41%	108 45%	228 39%	115 38%	222 42%
Not too serious	389 47%	47 43%	84 59% AC	131 43%	96 48%	31 45%	123 47%	267 48%	184 47%	206 48%	11 38%	378 48%	101 42%	289 50% L	153 51%	237 45%
Not at all serious	42 5%	7 7%	5 4%	12 4%	15 8%	3 4%	10 4%	32 6%	15 4%	27 6%	0 -	42 5%	11 5%	31 5%	14 5%	28 5%
(DK/NS)	5 1%	0 -	1 1%	1 0	3 2%	0 -	1 0	4 1%	3 1%	2 0	1 3%	4 1%	1 0	4 1%	2 1%	3 1%
Summary																
Top2Box	385 47%	55 50% B	52 37%	158 52% BD	85 43%	34 51% B	128 49%	257 46%	192 49%	194 45%	17 59%	368 46%	128 53% M	258 44%	130 43%	256 49%
Low2Box	431 52%	54 50%	89 63% AC	143 47%	111 56%	33 49%	133 51%	299 53%	199 50%	233 54%	11 38%	420 53%	112 46%	320 55% L	167 56%	265 51%

Rotavirus Study (60-6056-01)

Q6_5. [Forceful vomiting with increased frequency] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Forceful vomiting with increased frequency																
Very serious	647 79%	87 79%	105 73%	241 80%	161 80%	53 79%	223 85% G	423 76%	316 80%	331 77%	25 86%	622 78%	195 81%	451 78%	228 76%	419 80%
Serious	152 18%	17 16%	33 23%	56 18%	32 16%	14 21%	30 12%	121 22% F	71 18%	81 19%	3 10%	149 19%	39 16%	113 19%	62 21%	90 17%
Not too serious	5 1%	1 1%	2 1%	2 1%	0 -	0 -	4 2% G	1 0	1 0	4 1%	0 -	5 1%	0 -	5 1%	2 1%	3 1%
Not at all serious	15 2%	4 4%	2 1%	3 1%	6 3%	0 -	4 2%	11 2%	4 1%	11 3%	0 -	15 2%	5 2%	10 2%	6 2%	9 2%
(DK/NS)	3 0	0 -	1 1%	1 0	1 1%	0 -	0 -	3 1%	1 0	2 0	1 3%	2 0	1 0	2 0	1 0	2 0
Summary																
Top2Box	799 97%	105 95%	138 96%	297 98%	193 96%	67 100%	254 97%	545 97%	387 98% I	411 96%	28 97%	770 97%	234 97%	565 97%	289 97%	509 97%
Low2Box	20 2%	5 5%	4 3%	5 2%	6 3%	0 -	8 3%	12 2%	5 1%	15 4% H	0 -	20 3%	5 2%	15 3%	8 3%	12 2%

Rotavirus Study (60-6056-01)

Q6_6. [Decreased activity] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Decreased activity																
Very serious	119	14	28	54	13	10	38	81	49	70	5	114	43	76	46	73
	15%	13%	19% D	18% D	7%	14% D	15%	14%	12%	16%	17%	14%	18%	13%	16%	14%
Serious	329	49	48	136	70	25	100	230	169	160	16	313	98	231	131	198
	40%	45%	34% 45% BD	35% BD	35%	38% BD	38%	41%	43%	37%	55%	39%	41%	40%	44%	38%
Not too serious	337	39	64	100	103	31	110	227	158	179	7	330	89	248	109	228
	41%	36%	45% C	33% AC	52% AC	47% C	42%	41%	40%	42%	24%	42%	37%	43%	37%	44% N
Not at all serious	32	7	2	10	12	1	14	18	16	16	0	32	9	23	10	22
	4%	7% B	1%	3%	6% B	1%	5%	3%	4%	4%	-	4%	4%	4%	3%	4%
(DK/NS)	4	0	1	2	1	0	0	4	1	3	1	3	1	3	2	2
	0	-	1%	1%	1%	-	-	1%	0	1%	3%	0	0	1%	1%	0
Summary																
Top2Box	448	64	76	191	83	35	138	310	218	231	21	427	141	307	177	271
	55%	58% D	53% D	63% D	42%	52%	53%	55%	55%	54%	73%	54%	59%	53%	59% O	52%
Low2Box	370	46	66	110	115	32	124	246	175	195	7	363	98	272	119	250
	45%	42%	46%	36%	58% ABC	48%	47%	44%	44%	45%	24%	46%	41%	47%	40%	48% N

Rotavirus Study (60-6056-01)

Q6_7. [Hives] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Hives																
Very serious	135 16%	15 14%	27 19% D	68 22% D	15 8%	10 16% D	47 18%	88 16%	57 14%	78 18%	7 24%	128 16%	42 17%	93 16%	48 16%	88 17%
Serious	359 44%	61 55% DE	62 43%	138 46%	74 37%	24 36%	112 43%	247 44%	181 46%	178 42%	13 45%	346 44%	107 45%	252 43%	140 47%	219 42%
Not too serious	277 34%	29 26%	45 32%	86 28%	88 44% ABC	29 43% AC	85 32%	192 34%	144 36%	133 31%	8 27%	269 34%	82 34%	195 34%	95 32%	182 35%
Not at all serious	22 3%	4 4%	4 3%	5 2%	8 4%	1 1%	9 3%	13 2%	5 1%	17 4% H	0 -	22 3%	4 2%	18 3%	5 2%	17 3%
(DK/NS)	29 4%	1 1%	5 4%	6 2%	14 7% AC	3 4%	9 3%	20 4%	7 2%	22 5% H	1 3%	28 4%	6 2%	23 4%	11 4%	18 3%
Summary																
Top2Box	494 60%	76 69% DE	88 62% D	206 68% DE	89 45%	35 52%	159 61%	335 60%	238 60%	256 60%	20 69%	474 60%	149 62%	346 59%	188 63%	307 59%
Low2Box	299 36%	33 30%	49 35%	91 30%	96 48% ABC	30 44% AC	94 36%	205 37%	149 38%	150 35%	8 27%	291 37%	86 36%	213 37%	100 33%	199 38%

Rotavirus Study (60-6056-01)

Q6_8. [Severe diarrhea] Here is a list of symptoms that could affect your child/ children. Thinking about your youngest child, please indicate how serious you think each symptom is —very serious, serious, not too serious, or not at all serious?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Severe diarrhea																
Very serious	617 75%	83 76%	107 75%	245 81% DE	134 67%	47 70%	207 79%	409 73%	304 77%	313 73%	21 73%	595 75%	193 80% M	424 73%	207 69%	410 78% N
Serious	174 21%	22 20%	32 22%	48 16% C	53 27% C	19 29% C	46 17%	129 23%	81 21%	93 22%	8 27%	166 21%	45 19%	130 22%	77 26% O	97 18%
Not too serious	13 2%	0 -	1 1%	6 2%	6 3%	0 -	3 1%	10 2%	4 1%	9 2%	0 -	13 2%	0 -	13 2% L	8 3%	5 1%
Not at all serious	14 2%	5 5% C	2 1%	3 1%	4 2%	0 -	5 2%	9 2%	3 1%	11 3% H	0 -	14 2%	3 1%	11 2%	4 1%	10 2%
(DK/NS)	4 0	0 -	1 1%	0 -	2 1%	1 1% C	1 0	3 1%	2 1%	2 0	0 -	4 0	0 -	4 1%	2 1%	2 0
Summary																
Top2Box	791 96%	105 95%	139 97%	294 97%	188 94%	66 99%	253 97%	538 96%	384 98% I	406 95%	29 100%	762 96%	237 99% M	554 95%	285 95%	506 97%
Low2Box	27 3%	5 5%	3 2%	9 3%	10 5%	0 -	8 3%	19 3%	7 2%	20 5% H	0 -	27 3%	3 1%	24 4% L	12 4%	15 3%

Rotavirus Study (60-6056-01)

Q7. Have you ever heard of a virus called rotavirus, which causes gastroenteritis ... ?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Yes	393	63	64	137	93	37	121	273	393	0	29	364	212	181	143	250
	48%	57%	45%	45%	47%	55%	46%	49%	100%	-	100%	46%	88%	31%	48%	48%
		C							I				M			
No	429	47	79	166	106	31	141	287	0	429	0	429	28	400	155	273
	52%	43%	55%	55%	53%	45%	54%	51%	-	100%	-	54%	12%	69%	52%	52%
		A							H				L			

Rotavirus Study (60-6056-01)

Q8. How familiar would you say you are with the rotavirus —very familiar, familiar, not too familiar, or not at all familiar?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: Ever heard about rotavirus Weighted	394	61	62	136	93	42	121	273	394	0	29	365	212	182	144	250
	393	63*	64*	137	93*	37*	121	273	393	-**	29**	364	212	181	143	250
Very familiar	35	3	7	18	4	3	9	26	35	0	5	30	32	3	22	13
	9%	5%	11%	13% D	4%	7%	7%	10%	9%	-	17%	8%	15% M	2%	15% O	5%
Familiar	126	19	14	53	29	10	44	82	126	0	8	118	82	44	45	81
	32%	31%	23%	39% B	31%	26%	37%	30%	32%	-	28%	32%	39% M	24%	31%	33%
Not too familiar	208	38	39	57	53	22	65	144	208	0	15	193	90	118	68	140
	53%	61% C	61% C	41%	57% C	60% C	54%	53%	53%	-	51%	53%	43% L	65% L	47%	56%
Not at all familiar	24	2	3	9	7	3	3	21	24	0	1	23	8	16	9	15
	6%	3%	5%	7%	8%	7%	2%	8% F	6%	-	3%	6%	4% L	9% L	6%	6%
Summary																
Top2Box	161	23	22	72	33	12	53	108	161	0	13	148	114	47	67	94
	41%	36%	34%	52% ABDE	35%	33%	44%	40%	41%	-	45%	41%	54% M	26%	47%	38%
Low2Box	232	40	42	66	60	24	68	164	232	0	16	216	98	134	77	156
	59%	64% C	66% C	48% C	65% C	67% C	56%	60%	59%	-	55%	59%	46% L	74% L	53%	62%

Rotavirus Study (60-6056-01)

Q9. Has your youngest child received vaccination ... for the rotavirus?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: Ever heard about rotavirus Weighted	394	61	62	136	93	42	121	273	394	0	29	365	212	182	144	250
	393	63*	64*	137	93*	37*	121	273	393	-**	29**	364	212	181	143	250
Yes	29 7%	9 15%	4 6%	9 7%	5 5%	2 5%	15 12% G	14 5%	29 7%	0 -	29 100%	0 -	21 10% M	8 4%	10 7%	19 8%
No	364 93%	53 85%	60 94%	128 93%	88 95%	35 95%	106 88%	258 95% F	364 93%	0 -	0 -	364 100%	191 90%	173 96% L	133 93%	231 92%

Rotavirus Study (60-6056-01)

Q10. [SUMMARY - TOP2BOX] Here is a list of conditions that could affect your child. Thinking about your youngest child, please indicate how serious you think each condition is-very serious, serious, not too serious or not at all serious

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Whooping Cough	746 91%	102 93%	129 91%	273 90%	178 89%	64 95%	240 92%	506 90%	361 92%	384 90%	29 100%	716 90%	226 94% M	519 89%	273 92%	472 90%
Measles	687 84%	95 87% D	119 83% D	265 88% D	149 74%	58 87% D	218 83%	469 84%	335 85%	352 82%	27 94%	659 83%	211 88% M	475 82%	248 83%	439 84%
Rotavirus	620 75%	82 75%	108 76%	239 79% D	140 70%	51 75%	201 77%	419 75%	322 82% I	298 70%	24 83%	596 75%	207 86% M	413 71%	222 74%	398 76%
Influenza	562 68%	82 75% D	107 75% D	241 80% D	81 41%	51 75% D	189 72%	373 67%	277 70%	285 66%	23 80%	539 68%	167 70%	395 68%	198 66%	364 70%
Chicken Pox	476 58%	79 72% BD	84 59% D	192 63% D	77 39%	44 65% D	161 61%	315 56%	226 57%	250 58%	24 83%	451 57%	150 63%	325 56%	175 59%	301 57%

Rotavirus Study (60-6056-01)

Q10. [SUMMARY - LOW2BOX] Here is a list of conditions that could affect your child. Thinking about your youngest child, please indicate how serious you think each condition is-very serious, serious, not too serious or not at all serious

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Chicken Pox	341 42%	31 28%	57 40%	108 36%	121 61%	24 35%	99 38%	242 43%	167 42%	175 41%	5 17%	336 42%	90 37%	251 43%	122 41%	220 42%
Influenza	248 30%	26 23%	33 23%	56 18%	117 59%	17 25%	68 26%	180 32%	117 30%	131 31%	6 20%	242 31%	72 30%	176 30%	98 33%	150 29%
Measles	125 15%	14 13%	22 15%	33 11%	47 24%	9 13%	39 15%	86 15%	57 14%	68 16%	2 6%	123 16%	28 12%	97 17%	47 16%	78 15%
Rotavirus	96 12%	13 12%	10 7%	25 8%	44 22%	3 4%	22 8%	73 13%	51 13%	44 10%	2 7%	94 12%	26 11%	70 12%	45 15%	50 10%
Whooping Cough	58 7%	6 6%	8 6%	24 8%	16 8%	3 5%	14 5%	44 8%	30 8%	28 7%	0 -	58 7%	13 5%	45 8%	20 7%	38 7%

Rotavirus Study (60-6056-01)

Q10_1. [Rotavirus] Here is a list of conditions that could affect your child. Thinking about your youngest child, please indicate how serious you think each condition is—very serious, serious, not too serious or not at all serious.

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Rotavirus																
Very serious	237 29%	29 26%	34 24%	105 35% BD	45 23%	24 36% BD	96 36% G	142 25%	116 29%	122 28%	14 49%	223 28%	92 38% M	145 25%	87 29%	150 29%
Serious	383 47%	53 49%	74 52%	134 44%	95 48%	26 39%	105 40%	278 50% F	206 52% I	177 41%	10 35%	373 47%	115 48%	268 46%	135 45%	248 47%
Not too serious	93 11%	13 12%	10 7%	23 8%	43 22% ABCE	3 4%	21 8%	71 13% F	50 13%	42 10%	2 7%	91 11%	26 11%	67 11%	44 15% O	48 9%
Not at all serious	3 0	0 -	0 -	2 1%	1 1%	0 -	1 0	2 0	1 0	2 0	0 -	3 0	0 -	3 1%	1 0	2 0
(DK/NS)	106 13%	14 13%	25 17% D	38 13%	15 8%	14 21% D	39 15%	67 12%	20 5%	86 20% H	3 9%	104 13%	7 3%	100 17% L	31 10%	75 14%
Summary																
Top2Box	620 75%	82 75%	108 76%	239 79% D	140 70%	51 75%	201 77%	419 75%	322 82% I	298 70%	24 83%	596 75%	207 86% M	413 71%	222 74%	398 76%
Low2Box	96 12%	13 12%	10 7%	25 8%	44 22% ABCE	3 4%	22 8%	73 13%	51 13%	44 10%	2 7%	94 12%	26 11%	70 12%	45 15% O	50 10%

Rotavirus Study (60-6056-01)

Q10_2. [Influenza] Here is a list of conditions that could affect your child. Thinking about your youngest child, please indicate how serious you think each condition is—very serious, serious, not too serious or not at all serious.

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Influenza																
Very serious	195	30	35	99	13	18	70	125	91	104	15	180	74	121	79	116
	24%	27% D	24% D	33% D	7%	27% D	27%	22%	23%	24%	52%	23%	31% M	21%	27%	22%
Serious	367	52	72	142	68	32	119	248	186	181	8	359	93	274	118	249
	45%	48% D	50% D	47% D	34%	48% D	45%	44%	47%	42%	28%	45%	39% L	47% L	40%	48% N
Not too serious	226	26	31	51	101	17	61	165	106	119	6	220	66	160	85	141
	27%	23%	22%	17%	51% ABCE	25%	23%	29%	27%	28%	20%	28%	27%	28%	28%	27%
Not at all serious	22	0	2	4	16	0	7	15	10	12	0	22	6	16	13	9
	3%	-	1%	1%	8% ABCE	-	3%	3%	3%	3%	-	3%	3%	3%	4% O	2%
(DK/NS)	12	2	3	6	1	0	5	7	0	12	0	12	1	11	3	9
	1%	2%	2%	2%	1%	-	2%	1%	-	3% H	-	2%	0	2%	1%	2%
Summary																
Top2Box	562	82	107	241	81	51	189	373	277	285	23	539	167	395	198	364
	68%	75% D	75% D	80% D	41%	75% D	72%	67%	70%	66%	80%	68%	70%	68%	66%	70%
Low2Box	248	26	33	56	117	17	68	180	117	131	6	242	72	176	98	150
	30%	23%	23%	18%	59% ABCE	25%	26%	32%	30%	31%	20%	31%	30%	30%	33%	29%

Rotavirus Study (60-6056-01)

Q10_3. [Whooping Cough] Here is a list of conditions that could affect your child. Thinking about your youngest child, please indicate how serious you think each condition is—very serious, serious, not too serious or not at all serious.

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Whooping Cough																
Very serious	405 49%	56 51%	68 47%	151 50%	96 48%	33 49%	144 55% G	261 47%	190 48%	215 50%	18 63%	387 49%	135 56% M	270 46%	147 49%	258 49%
Serious	341 41%	45 41%	62 43%	122 40%	81 41%	31 45%	96 37%	244 44%	171 43%	170 40%	11 37%	330 42%	91 38%	250 43%	126 42%	214 41%
Not too serious	56 7%	6 6%	8 6%	23 8%	15 8%	3 5%	14 5%	42 8%	30 8%	26 6%	0 -	56 7%	13 5%	43 7%	19 6%	37 7%
Not at all serious	2 0	0 -	0 -	1 0	1 1%	0 -	0 -	2 0	0 -	2 0	0 -	2 0	0 -	2 0	1 0	1 0
(DK/NS)	18 2%	2 2%	5 4%	5 2%	6 3%	0 -	8 3%	10 2%	2 1%	16 4% H	0 -	18 2%	1 0	17 3% L	5 2%	13 3%
Summary																
Top2Box	746 91%	102 93%	129 91%	273 90%	178 89%	64 95%	240 92%	506 90%	361 92%	384 90%	29 100%	716 90%	226 94% M	519 89%	273 92%	472 90%
Low2Box	58 7%	6 6%	8 6%	24 8%	16 8%	3 5%	14 5%	44 8%	30 8%	28 7%	0 -	58 7%	13 5%	45 8%	20 7%	38 7%

Rotavirus Study (60-6056-01)

Q10_4. [Chicken Pox] Here is a list of conditions that could affect your child. Thinking about your youngest child, please indicate how serious you think each condition is—very serious, serious, not too serious or not at all serious.

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Chicken Pox																
Very serious	155 19%	32 29% DE	31 22% D	66 22% D	16 8%	10 16%	60 23% G	94 17%	76 19%	79 18%	12 42%	142 18%	60 25% M	95 16%	57 19%	98 19%
Serious	321 39%	47 43% D	53 37% D	126 42% D	61 31%	33 49% D	101 38%	221 39%	150 38%	171 40%	12 41%	309 39%	91 38%	230 40%	118 40%	203 39%
Not too serious	315 38%	29 26%	52 37%	99 33%	112 56% ABCE	23 34%	94 36%	221 39%	152 39%	163 38%	5 17%	310 39%	83 34%	232 40%	111 37%	204 39%
Not at all serious	26 3%	2 2%	5 4%	9 3%	9 5%	1 1%	5 2%	21 4%	14 4%	12 3%	0 -	26 3%	7 3%	19 3%	11 4%	15 3%
(DK/NS)	5 1%	0 -	1 1%	3 1%	1 1%	0 -	2 1%	3 1%	1 0	4 1%	0 -	5 1%	0 -	5 1%	2 1%	3 1%
Summary																
Top2Box	476 58%	79 72% BD	84 59% D	192 63% D	77 39%	44 65% D	161 61%	315 56%	226 57%	250 58%	24 83%	451 57%	150 63%	325 56%	175 59%	301 57%
Low2Box	341 42%	31 28%	57 40% A	108 36%	121 61% ABCE	24 35%	99 38%	242 43%	167 42%	175 41%	5 17%	336 42%	90 37%	251 43%	122 41%	220 42%

Rotavirus Study (60-6056-01)

Q10_5. [Measles] Here is a list of conditions that could affect your child. Thinking about your youngest child, please indicate how serious you think each condition is—very serious, serious, not too serious or not at all serious.

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Measles																
Very serious	309	55	47	131	50	25	109	200	149	160	18	291	109	200	115	194
	38%	50% BD	33%	43% BD	25%	38% D	42%	36%	38%	37%	63%	37%	46% M	34%	39%	37%
Serious	378	40	72	134	98	33	109	269	185	192	9	369	102	276	132	245
	46%	36%	50% A	44%	49% A	49%	41%	48%	47%	45%	31%	46%	42%	47%	44%	47%
Not too serious	118	14	22	30	43	9	37	81	55	63	2	116	26	92	43	75
	14%	13%	15%	10%	22% C	13%	14%	14%	14%	15%	6%	15%	11%	16%	14%	14%
Not at all serious	7	0	0	3	4	0	2	5	2	5	0	7	2	5	4	3
	1%	-	-	1%	2%	-	1%	1%	1%	1%	-	1%	1%	1%	1%	1%
(DK/NS)	10	0	2	4	4	0	5	5	2	8	0	10	1	9	4	6
	1%	-	1%	1%	2%	-	2%	1%	1%	2%	-	1%	0	2%	1%	1%
Summary																
Top2Box	687	95	119	265	149	58	218	469	335	352	27	659	211	475	248	439
	84%	87% D	83% D	88% D	74%	87% D	83%	84%	85%	82%	94%	83%	88% M	82%	83%	84%
Low2Box	125	14	22	33	47	9	39	86	57	68	2	123	28	97	47	78
	15%	13%	15%	11%	24% AC	13%	15%	15%	14%	16%	6%	16%	12%	17%	16%	15%

Rotavirus Study (60-6056-01)

Q11. [SUMMARY - YES] Here are the same conditions. For each, please indicate if you have seen, read, or heard anything about it in the past 6 months.

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
Base: All respondents Weighted		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Influenza	655	91	103	241	164	57	211	444	343	313	27	628	220	435	252	404
	80%	83% B	72%	80%	82% B	84% B	81%	79%	87% I	73%	93%	79%	92% M	75%	84% O	77%
Chicken Pox	432	70	65	166	99	33	148	284	205	228	19	413	148	284	168	264
	53%	64% BD	45%	55%	50%	49%	56%	51%	52%	53%	66%	52%	62% M	49%	56%	50%
Whooping Cough	279	36	44	129	51	18	106	173	132	146	19	259	109	169	113	165
	34%	33%	31%	43% BDE	26%	27%	41% G	31%	34%	34%	66%	33%	46% M	29%	38%	32%
Rotavirus	240	36	35	78	74	17	84	156	212	28	21	219	240	0	104	136
	29%	33%	24%	26%	37% BC	26%	32%	28%	54% I	7%	73%	28%	100% M	-	35% O	26%
Measles	238	39	39	85	58	17	95	143	110	127	20	217	93	144	85	152
	29%	36%	27%	28%	29%	25%	36% G	25%	28%	30%	69%	27%	39% M	25%	29%	29%

Rotavirus Study (60-6056-01)

Q12. From what sources have you heard about the rotavirus ... ?

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: Ever heard about rotavirus	394	61	62	136	93	42	121	273	394	0	29	365	212	182	144	250
Weighted	393	63*	64*	137	93*	37*	121	273	393	-**	29**	364	212	181	143	250
Television	153 39%	19 31%	21 32%	60 43%	41 44%	12 33%	51 42%	102 37%	153 39%	0 -	8 27%	145 40%	104 49% M	49 27%	59 41%	94 37%
Friends	122 31%	12 20%	21 32%	46 34% A	28 30%	15 40% A	44 36%	79 29%	122 31%	0 -	9 31%	113 31%	79 37% M	43 24%	52 36%	70 28%
Brochures/ pamphlets/literature available in doctor's office	117 30%	24 38% DE	17 27%	48 35% D	20 22%	7 19%	44 37% G	72 26%	117 30%	0 -	16 56%	100 28%	78 37% M	39 21%	41 29%	75 30%
Magazines	100 25%	19 31% D	13 21%	46 34% D	12 13%	9 24%	31 26%	69 25%	100 25%	0 -	6 21%	94 26%	65 31% M	35 19%	33 23%	67 27%
Internet	81 21%	10 16%	11 18%	34 25%	17 18%	8 21%	30 25%	51 19%	81 21%	0 -	3 10%	78 21%	51 24%	30 17%	36 25%	45 18%
Parent/ family member	69 18%	5 8%	5 8%	26 19% B	23 25% AB	10 26% AB	26 21%	43 16%	69 18%	0 -	5 17%	64 18%	48 22% M	22 12%	31 21%	39 15%
General practitioner / Family medicine doctor	68 17%	10 16%	14 23% D	26 19%	9 10%	8 21%	20 16%	48 18%	68 17%	0 -	11 39%	57 16%	51 24% M	17 9%	30 21%	38 15%
Newspaper articles	63 16%	8 13%	6 10%	18 13%	23 25% BC	7 19%	21 17%	42 15%	63 16%	0 -	3 10%	60 16%	40 19%	23 13%	23 16%	39 16%
Nurse	52 13%	10 16%	8 13%	13 10%	16 17%	4 12%	18 15%	34 13%	52 13%	0 -	11 38%	41 11%	39 18% M	13 7%	22 15%	30 12%
Pediatrician	42 11%	2 3%	5 8%	18 13% A	16 17% AE	1 2%	16 13%	26 10%	42 11%	0 -	10 35%	32 9%	34 16% M	8 4%	18 13%	24 10%
Emergency room physician	38 10%	3 5%	3 5%	14 10%	12 13%	5 14%	11 9%	27 10%	38 10%	0 -	2 7%	36 10%	23 11%	15 8%	17 12%	20 8%
Books	37 9%	4 7%	4 6%	18 13%	7 8%	3 10%	14 11%	23 8%	37 9%	0 -	6 21%	31 8%	25 12%	12 7%	14 10%	23 9%

Rotavirus Study (60-6056-01)

Q12. From what sources have you heard about the rotavirus ... ?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Radio	34 9%	1 2%	4 6%	11 8%	14 15% A	3 10%	12 10%	22 8%	34 9%	0 -	1 3%	33 9%	27 13% M	7 4%	12 8%	22 9%
Pharmacist	20 5%	1 2%	0 -	5 4%	13 14% ABCE	1 2%	9 7%	11 4%	20 5%	0 -	2 7%	18 5%	13 6%	7 4%	11 8%	9 4%
Teacher	10 2%	1 2%	0 -	6 4%	1 1%	2 5%	3 3%	7 2%	10 2%	0 -	0 -	10 3%	7 3%	3 2%	4 3%	6 2%
Knew someone who had contracted it	6 2%	0 -	1 2%	4 3%	1 1%	0 -	2 2%	4 1%	6 2%	0 -	0 -	6 2%	5 2%	1 1%	6 4% O	0 -
At work	4 1%	1 2%	0 -	3 2%	0 -	0 -	1 1%	3 1%	4 1%	0 -	0 -	4 1%	2 1%	2 1%	0 -	4 2%
School	2 1%	1 2%	0 -	1 1%	0 -	0 -	0 -	2 1%	2 1%	0 -	0 -	2 1%	0 -	2 1%	0 -	2 1%
Other	5 1%	1 2%	0 -	2 1%	2 2%	0 -	3 3%	2 1%	5 1%	0 -	1 4%	4 1%	3 1%	2 1%	2 1%	3 1%
(DK/NS)	9 2%	1 2%	4 6%	2 1%	1 1%	1 2%	2 2%	7 3%	9 2%	0 -	1 3%	8 2%	0 -	9 5% L	1 1%	8 3%

Rotavirus Study (60-6056-01)

Q13. Please describe the information you saw or heard about the rotavirus ... ?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: Ever heard about rotavirus	394	61	62	136	93	42	121	273	394	0	29	365	212	182	144	250
Weighted	393	63*	64*	137	93*	37*	121	273	393	-**	29**	364	212	181	143	250
Causes diarrhea	60 15%	6 10%	7 11%	26 19%	13 14%	7 19%	22 18%	38 14%	60 15%	0 -	5 17%	55 15%	39 18%	21 11%	26 18%	34 13%
That it's especially affects younger children/ children under 5 years of age	43 11%	6 10%	8 13%	13 10%	12 13%	3 10%	15 13%	28 10%	43 11%	0 -	5 17%	38 10%	30 14% M	13 7%	15 10%	28 11%
Easily transmitted/ very contagious	41 10%	6 10%	3 5%	16 12%	11 12%	4 12%	15 12%	26 9%	41 10%	0 -	2 7%	39 11%	33 15% M	8 4%	13 9%	28 11%
That there is a vaccine for the rotavirus/ saw a vaccine ad	38 10%	6 10%	3 5%	16 12%	10 11%	3 7%	11 9%	27 10%	38 10%	0 -	3 10%	35 10%	32 15% M	6 3%	18 12%	20 8%
The symptoms (unspecified)	31 8%	7 11%	3 5%	11 8%	6 6%	3 10%	18 15% G	13 5%	31 8%	0 -	3 10%	28 8%	19 9%	12 7%	10 7%	21 8%
Family / friends had the virus/ were hospitalized because of it	30 8%	7 11%	5 8%	13 10%	4 4%	1 2%	8 7%	22 8%	30 8%	0 -	2 7%	28 8%	17 8%	13 7%	22 16% O	8 3%
Causes dehydration	28 7%	4 7%	3 5%	6 4%	11 12% C	3 10%	10 8%	18 7%	28 7%	0 -	3 10%	25 7%	22 10% M	6 3%	9 6%	19 8%
Causes vomiting	26 7%	2 3%	2 3%	11 8%	8 9%	3 7%	11 9%	15 5%	26 7%	0 -	3 10%	23 6%	17 8%	9 5%	11 8%	15 6%
Wash hands thoroughly to avoid spreading	19 5%	5 8%	3 5%	4 3%	7 8%	0 -	7 6%	12 4%	19 5%	0 -	3 11%	16 4%	17 8% M	2 1%	6 4%	13 5%
That the rotavirus could be very dangerous (sometimes cause even death) if not treated	19 5%	1 2%	2 3%	6 4%	9 10% A	1 2%	5 4%	14 5%	19 5%	0 -	0 -	19 5%	11 5%	8 4%	10 7%	9 4%
That there is an outbreak of this illness this winter	18 5%	1 2%	2 3%	7 5%	7 8%	1 2%	6 5%	12 4%	18 5%	0 -	0 -	18 5%	11 5%	7 4%	6 4%	12 5%

Rotavirus Study (60-6056-01)

Q13. Please describe the information you saw or heard about the rotavirus ...?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Just read about it in a print ad	18 4%	2 3%	3 5% D	9 7% D	0 -	3 10% D	6 5%	12 4%	18 4%	0 -	0 -	18 5%	10 5%	8 4%	8 5%	10 4%
Saw a Lysol commercial	17 4%	4 7%	1 2%	10 7% D	1 1%	1 2%	6 5%	11 4%	17 4%	0 -	2 7%	15 4%	10 5%	7 4%	8 6%	9 4%
Gastroenteritis	15 4%	0 -	1 2%	1 1%	12 13% ABC	1 2%	7 6%	8 3%	15 4%	0 -	1 3%	14 4%	10 5%	5 3%	4 3%	11 4%
Upset stomach	13 3%	2 3%	4 6%	6 4%	1 1%	0 -	4 3%	9 3%	13 3%	0 -	2 7%	11 3%	10 5%	3 2%	5 4%	8 3%
Lasts 3-4 days	11 3%	1 2%	3 5%	2 1%	3 3%	2 5%	2 2%	9 3%	11 3%	0 -	1 4%	10 3%	8 4%	3 2%	6 4%	5 2%
Causes fever	9 2%	0 -	1 2%	4 3%	3 3%	1 2%	4 3%	5 2%	9 2%	0 -	1 3%	8 2%	5 2%	4 2%	4 3%	5 2%
Heard it in the news	6 2%	1 2%	1 2%	3 2%	1 1%	0 -	1 1%	5 2%	6 2%	0 -	0 -	6 2%	4 2%	2 1%	3 2%	3 1%
How you can contract it	6 1%	2 3% C	0 -	0 -	2 2%	2 5% C	2 2%	4 1%	6 1%	0 -	1 4%	5 1%	1 0	5 3%	1 1%	5 2%
Just remember the name	22 6%	8 13% CD	5 8%	6 4%	2 2%	1 2%	9 8%	13 5%	22 6%	0 -	0 -	22 6%	3 1%	19 11% L	6 4%	16 7%
None/ haven't heard anything	11 3%	2 3%	4 6%	3 2%	1 1%	1 2%	2 2%	9 3%	11 3%	0 -	0 -	11 3%	1 0	10 6% L	3 2%	8 3%
Other	40 10%	4 7%	10 16%	12 9%	7 8%	7 19% D	13 11%	27 10%	40 10%	0 -	3 10%	37 10%	26 12%	15 8%	17 12%	24 10%
Dk/ ns	65 17%	8 13%	13 21%	25 18%	13 14%	5 14%	9 7%	56 21% F	65 17%	0 -	8 27%	57 16%	22 10%	43 24% L	16 11%	49 20% N

Rotavirus Study (60-6056-01)

Q14. Has your youngest child ever experienced five or more episodes of profuse, watery diarrhea in one day?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Yes	183 22%	30 27%	26 18%	64 21%	49 25%	15 22%	66 25%	117 21%	90 23%	93 22%	8 27%	175 22%	63 26%	120 21%	183 61% O	0 -
No	639 78%	80 73%	117 82%	239 79%	151 75%	52 78%	196 75%	443 79%	304 77%	335 78%	21 73%	618 78%	177 74%	462 79%	116 39%	523 100% N

Rotavirus Study (60-6056-01)

Q15. How many times has your youngest child experienced this in his/ her lifetime?

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: The youngest child who has experienced profuse, watery diarrhea Weighted	183	29	25	63	49	17	66	117	90	93	8	175	63	120	183	0
	183	30**	26**	64*	49*	15**	66*	117	90*	93*	8**	175	63*	120	183	-**
	1	98	15	11	35	28	8	37	61	52	46	4	94	32	66	98
		54%	52%	44%	56%	57%	53%	57%	52%	58%	49%	51%	54%	51%	55%	54%
	2	48	5	7	14	16	5	15	33	22	26	2	46	17	31	48
		26%	17%	28%	22%	33%	35%	22%	28%	24%	28%	24%	26%	27%	26%	26%
	3	11	3	2	4	2	0	3	8	7	4	2	9	7	4	11
		6%	10%	8%	6%	4%	-	5%	7%	8%	4%	26%	5%	11% M	3%	6%
4	7	1	2	4	0	0	3	4	1	6	0	7	1	6	7	0
		4%	3%	8%	6%	-	5%	3%	1%	7%	-	4%	2%	5%	4%	-
5	3	3	0	0	0	0	2	1	2	1	0	3	2	1	3	0
		2%	10%	-	-	-	3%	1%	2%	1%	-	2%	3%	1%	2%	-
6+	16	2	3	6	3	2	6	10	6	10	0	16	4	12	16	0
		9%	7%	12%	10%	12%	9%	9%	7%	11%	-	9%	6%	10%	9%	-
Summary																
Mean	3.1	2.7	2.8	3.1	3.7	2.1	2.9	3.2	2.3	3.9	1.7	3.1	2.8	3.3	3.1	0
Std Dev	7.7	3.7	3.3	6.0	12.8	2.0	5.4	8.8	3.0	10.4	0.9	7.9	5.3	8.7	7.7	0
Std Dev	0.6	0.7	0.7	0.8	1.8	0.5	0.7	0.8	0.3	1.1	0.3	0.6	0.7	0.8	0.6	0
Median	1.0	1.0	1.2	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	0

Rotavirus Study (60-6056-01)

Q16. Has/ have your other child/ children ever experienced five or more episodes of profuse, watery diarrhea in one day?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: Those who have 2 or more children Weighted	560	69	96	213	130	52	0	560	273	287	14	546	156	404	233	327
	560	71*	99*	215	130	45*	-**	560	273	287	14**	546	156	404	233	327
Yes	197	28	30	69	54	17	0	197	104	93	5	192	74	123	197	0
	35%	39%	30%	32%	42%	37%	-	35%	38%	32%	36%	35%	47% M	30%	85% O	-
No	363	43	69	146	76	29	0	363	169	194	9	354	82	281	36	327
	65%	61%	70%	68%	58%	63%	-	65%	62%	68%	64%	65%	53% L	70% L	15% N	100% N

Rotavirus Study (60-6056-01)

Q17. Thinking about the last time your youngest child had five or more episodes of profuse, watery diarrhea in one day, did you ...?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: Respondents whose youngest child experienced 5+ episodes of diarrhea in one day	183	29	25	63	49	17	66	117	90	93	8	175	63	120	183	0
Weighted	183	30**	26**	64*	49*	15**	66*	117	90*	93*	8**	175	63*	120	183	-**
Treat it at home	105 57%	22 72%	12 48%	35 56%	28 57%	8 53%	33 50%	72 62%	53 59%	52 56%	7 87%	98 56%	34 54%	71 59%	105 57%	0 -
Take your child to a doctor	80 44%	13 45%	13 52%	27 43%	19 39%	7 47%	33 50%	47 40%	35 39%	45 48%	1 11%	79 45%	33 52%	47 39%	80 44%	0 -
Take your child to a hospital emergency room	37 20%	4 14%	4 16%	10 16%	14 29%	4 29%	15 23%	22 19%	24 27% I	13 14%	1 13%	36 20%	16 25%	21 17%	37 20%	0 -
Call the health line	2 1%	1 3%	0 -	1 2%	0 -	0 -	0 -	2 2%	1 1%	1 1%	0 -	2 1%	0 -	2 2%	2 1%	0 -
Other	1 1%	0 -	0 -	0 -	1 2%	0 -	1 2%	0 -	0 -	1 1%	0 -	1 1%	0 -	1 1%	1 1%	0 -

Rotavirus Study (60-6056-01)

Q18. Did your child stay overnight at a hospital ... ?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: Those who took their youngest child to the hospital emergency room	37	4	4	10	14	5	15	22	24	13	1	36	16	21	37	0
	37*	4**	4**	10**	14**	4**	15**	22**	24**	13**	1**	36*	16**	21**	37*	-**
Weighted																
Yes	14 39%	3 75%	0 -	4 40%	7 50%	0 -	6 41%	8 37%	10 42%	4 32%	1 100%	13 37%	7 44%	7 34%	14 39%	0 -
No	23 61%	1 25%	4 100%	6 60%	7 50%	4 100%	9 59%	14 63%	14 58%	9 68%	0 -	23 63%	9 56%	14 66%	23 61%	0 -

Rotavirus Study (60-6056-01)

Q19. Has your youngest child or any of your children ever stayed for at least one night at a hospital because of multiple episodes of profuse and watery diarrhea?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: Respondents whose child has/ children have experienced 5+ episodes of diarrhea in a day Weighted	299	44	45	99	80	31	66	233	144	155	10	289	104	195	299	0
	299	45*	46*	100*	80*	27*	66*	233	143	155	10**	289	104	194	299	-**
	42 14%	8 18%	6 13%	12 12%	15 19% E	1 3%	8 12%	34 15%	24 17%	18 12%	2 20%	40 14%	20 19%	22 11%	42 14%	0 -
Yes																
No	256 86%	37 82%	40 87%	88 88%	65 81%	26 97% D	58 88%	198 85%	119 83%	137 88%	8 80%	248 86%	84 81%	172 89%	256 86%	0 -

Rotavirus Study (60-6056-01)

Q20. How well do you feel you managed the last incidence of profuse and watery diarrhea with your youngest child?

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: Treated youngest child at home Weighted																
	105	21	12	35	28	9	33	72	53	52	7	98	34	71	105	0
	105	22**	12**	35*	28**	8**	33*	72*	53*	52*	7**	98*	34*	71*	105	-**
Very well	53	10	8	13	18	3	12	41	24	29	4	49	16	37	53	0
	51%	48%	67%	37%	64%	44%	36%	57%	46%	56%	58%	50%	48%	52%	51%	-
Well	48	11	4	20	8	4	19	29	25	23	3	45	17	31	48	0
	46%	52%	33%	57%	29%	56%	57%	40%	47%	44%	42%	46%	49%	44%	46%	-
Not very well	1	0	0	1	0	0	0	1	1	0	0	1	1	0	1	0
	1%	-	-	3%	-	-	-	1%	2%	-	-	1%	3%	-	1%	-
Not well at all	1	0	0	0	1	0	1	0	1	0	0	1	0	1	1	0
	1%	-	-	-	4%	-	3%	-	2%	-	-	1%	-	1%	1%	-
(DK/NS)	2	0	0	1	1	0	1	1	2	0	0	2	0	2	2	0
	2%	-	-	3%	4%	-	3%	1%	4%	-	-	2%	-	3%	2%	-
Summary																
Top2Box	101	22	12	33	26	8	31	70	49	52	7	94	33	68	101	0
	96%	100%	100%	94%	93%	100%	94%	97%	92%	100% H	100%	96%	97%	96%	96%	-
Low2Box	2	0	0	1	1	0	1	1	2	0	0	2	1	1	2	0
	2%	-	-	3%	4%	-	3%	1%	4%	-	-	2%	3%	1%	2%	-

Rotavirus Study (60-6056-01)

Q21. The first time your youngest child experienced multiple episodes of profuse, watery diarrhea, how worried were you—very worried, worried, not too worried or not at all worried?

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: Respondents whose child has/ children have experienced 5+ episodes of diarrhea in a day	299	44	45	99	80	31	66	233	144	155	10	289	104	195	299	0
Weighted	299	45*	46*	100*	80*	27*	66*	233	143	155	10**	289	104	194	299	-**
Very worried	102 34%	21 45% E	13 29%	39 39%	23 29%	6 23%	33 50% G	69 30%	50 35%	52 34%	5 49%	97 34%	41 40%	61 31%	102 34%	0 -
Worried	134 45%	17 39%	19 42%	41 41%	41 51%	15 55%	28 42%	107 46%	66 46%	68 44%	4 41%	130 45%	43 41%	92 47%	134 45%	0 -
Not too worried	50 17%	6 14%	11 24%	15 15%	12 15%	5 19%	4 6%	46 20% F	21 14%	29 19%	1 10%	49 17%	16 15%	34 17%	50 17%	0 -
Not at all worried	12 4%	1 2%	2 4%	4 4%	4 5%	1 3%	1 2%	11 5%	6 4%	6 4%	0 -	12 4%	4 4%	8 4%	12 4%	0 -
Summary																
Top2Box	237 79%	38 84%	33 71%	81 81%	64 80%	21 77%	61 92% G	176 76%	117 81%	120 77%	9 90%	228 79%	84 81%	153 79%	237 79%	0 -
Low2Box	62 21%	7 16%	13 29%	19 19%	16 20%	6 23%	5 8%	57 24% F	27 19%	35 23%	1 10%	61 21%	20 19%	42 21%	62 21%	0 -

Rotavirus Study (60-6056-01)

Q22. How serious do you find this condition to be?

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: Respondents whose child has/ children have experienced 5+ episodes of diarrhea in a day	299	44	45	99	80	31	66	233	144	155	10	289	104	195	299	0
Weighted	299	45*	46*	100*	80*	27*	66*	233	143	155	10**	289	104	194	299	-**
Very serious	88 30%	17 39% DE	9 20%	45 45% BDE	12 15%	4 16%	22 34%	66 28%	38 27%	50 32%	3 30%	85 30%	33 32%	55 28%	88 30%	0 -
Serious	159 53%	23 50%	30 64% C	40 40%	47 59% C	19 71% C	34 51%	125 54%	82 58%	77 49%	6 60%	153 53%	58 56%	101 52%	159 53%	0 -
Not too serious	47 16%	5 11%	6 13%	13 13%	19 24%	3 13%	9 14%	38 16%	22 15%	25 16%	1 10%	46 16%	13 12%	34 17%	47 16%	0 -
Not at all serious	4 1%	0 -	1 2%	1 1%	2 3%	0 -	1 2%	3 1%	1 1%	3 2%	0 -	4 1%	0 -	4 2%	4 1%	0 -
Summary																
Top2Box	248 83%	40 89%	39 84%	86 86% D	59 74%	24 87%	56 85%	192 82%	121 84%	127 82%	9 90%	239 83%	91 88%	156 80%	248 83%	0 -
Low2Box	51 17%	5 11%	7 16%	14 14%	21 26% C	3 13%	10 15%	41 18%	23 16%	28 18%	1 10%	50 17%	13 12%	38 20%	51 17%	0 -

Rotavirus Study (60-6056-01)

Q23. To the best of your knowledge, which, if any, of the following are signs of dehydration among infants?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Dry mouth	639 78%	82 75%	103 72%	233 77%	174 87%	47 70%	207 79%	431 77%	310 79%	329 77%	25 87%	613 77%	190 79%	448 77%	242 81%	397 76%
A rash	61 7%	8 7%	12 9%	25 8%	10 5%	5 8%	25 10%	36 6%	28 7%	33 8%	4 13%	57 7%	21 9%	40 7%	20 7%	41 8%
A cough	72 9%	11 10%	13 9%	26 9%	17 9%	4 6%	29 11%	43 8%	29 7%	43 10%	3 10%	69 9%	27 11%	45 8%	27 9%	45 9%
Sunken eyes and soft spots on your child's head	571 70%	85 78%	105 73%	216 71%	126 63%	39 58%	168 64%	404 72%	306 78%	265 62%	20 69%	551 70%	185 77%	387 67%	218 73%	353 67%
Lethargy	697 85%	101 92%	125 88%	256 85%	158 79%	58 86%	216 82%	481 86%	355 90%	343 80%	24 83%	673 85%	215 90%	482 83%	256 86%	441 84%
Fewer diaper changes	728 89%	94 86%	128 90%	272 90%	170 85%	63 94%	224 85%	504 90%	352 90%	375 88%	25 86%	702 89%	212 88%	515 89%	268 90%	459 88%
Runny nose	28 3%	3 3%	3 2%	15 5%	3 2%	3 5%	12 5%	16 3%	7 2%	21 5%	1 3%	27 3%	8 3%	20 3%	12 4%	16 3%
None of the above	7 1%	0 -	2 1%	2 1%	3 2%	0 -	4 2%	3 1%	2 1%	5 1%	1 4%	6 1%	1 0	6 1%	1 0	6 1%
(DK/NS)	14 2%	1 1%	4 3%	6 2%	2 1%	1 1%	5 2%	9 2%	5 1%	9 2%	1 3%	13 2%	3 1%	11 2%	4 1%	10 2%

Rotavirus Study (60-6056-01)

Q24. Has your youngest child ever been diagnosed by a doctor with viral diarrhea or rotavirus?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Yes	46 6%	5 5%	6 4%	15 5%	19 10% CE	1 1%	17 7%	29 5%	29 7% I	17 4%	1 3%	45 6%	28 12% M	18 3%	34 11% O	12 2%
No	755 92%	103 93%	133 94%	278 92%	177 88%	64 96%	237 90%	518 92%	354 90%	400 93%	26 90%	728 92%	206 86% L	549 94%	247 83% N	507 97% N
(DK/NS)	21 3%	2 2%	3 2%	10 3%	4 2%	2 3%	8 3%	13 2%	10 3%	11 3%	2 6%	19 2%	6 2%	15 3%	17 6% O	4 1%

Rotavirus Study (60-6056-01)

Q25. Has/ have your other child/ children ever been diagnosed by a doctor with viral diarrhea or rotavirus?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: Those who have 2 or more children Weighted	560	69	96	213	130	52	0	560	273	287	14	546	156	404	233	327
	560	71*	99*	215	130	45*	-**	560	273	287	14**	546	156	404	233	327
Yes	58	7	8	18	21	3	0	58	41	17	3	55	34	24	52	6
	10%	10%	8%	8%	16%	8%	-	10%	15%	6%	21%	10%	22%	6%	22%	2%
No	488	64	88	188	109	39	0	488	224	264	10	478	119	369	172	317
	87%	90%	90%	87%	84%	87%	-	87%	82%	92%	71%	88%	76%	91%	74%	97%
(DK/NS)	14	0	2	9	0	3	0	14	8	6	1	13	3	11	9	5
	2%	-	2%	4%	-	6%	-	2%	3%	2%	7%	2%	2%	3%	4%	2%

Rotavirus Study (60-6056-01)

Q26. Which, if any, of the following are possible symptoms or consequences of viral diarrhea or rotavirus?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: Ever heard about rotavirus	394	61	62	136	93	42	121	273	394	0	29	365	212	182	144	250
Weighted	393	63*	64*	137	93*	37*	121	273	393	-**	29**	364	212	181	143	250
Cough	26 7%	6 10% D	4 6%	13 10% D	2 2%	1 2%	8 7%	18 7%	26 7%	0 -	4 14%	22 6%	22 10% M	4 2%	12 9%	14 6%
Frequent episodes of watery diarrhea	336 85%	57 92% B	47 74%	122 89% B	78 84%	31 83%	100 83%	236 86%	336 85%	0 -	23 80%	312 86%	189 89% M	147 81%	125 87%	210 84%
Rash	72 18%	12 20% D	9 15%	36 26% D	5 5%	10 26% D	21 17%	52 19%	72 18%	0 -	7 24%	66 18%	41 19%	32 17%	28 19%	45 18%
Forceful vomiting with increased frequency	240 61%	33 52%	30 47%	84 61%	72 77% ABCE	22 60%	75 62%	166 61%	240 61%	0 -	19 66%	221 61%	161 76% M	80 44%	96 67%	145 58%
Headache	162 41%	27 43%	22 34%	52 38%	43 46%	18 50%	44 36%	119 44%	162 41%	0 -	11 38%	151 41%	97 46%	65 36%	64 45%	98 39%
None of the above	1 0	1 2%	0 -	0 -	0 -	0 -	0 -	1 0	1 0	0 -	0 -	1 0	0 -	1 1%	1 1%	0 -
(DK/NS)	53 13%	4 7%	15 24% AC	15 11%	12 13%	6 17%	20 16%	33 12%	53 13%	0 -	6 20%	47 13%	20 9%	33 18% L	14 10%	39 16%

Rotavirus Study (60-6056-01)

Q27. [SUMMARY - TRUE] Now, please indicate if you think the following statements about rotavirus or viral diarrhea are true or false.

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: Ever heard about rotavirus Weighted	394	61	62	136	93	42	121	273	394	0	29	365	212	182	144	250
	393	63*	64*	137	93*	37*	121	273	393	-**	29**	364	212	181	143	250
Severe rotavirus quickly leads to dehydration	349	56	56	120	86	30	105	244	349	0	22	327	192	157	130	219
	89%	90%	89%	88%	92% E	81%	87%	89%	89%	-	77%	90%	91%	87%	90%	88%
Rotavirus can be very serious and unpredictable	299	46	42	114	69	28	96	203	299	0	19	280	167	133	109	190
	76%	74%	66%	83% B	74%	76%	80%	75%	76%	-	66%	77%	79%	73%	76%	76%
Rotavirus is contagious	283	43	34	96	85	24	87	196	283	0	16	266	165	118	110	173
	72%	69%	53%	70% B	91% ABCE	67%	72%	72%	72%	-	56%	73%	78% M	65%	76%	69%
Rotavirus is the most common cause of diarrhea in young children	173	25	22	63	51	13	53	120	173	0	14	159	111	62	69	104
	44%	39%	34%	46%	55% BE	36%	44%	44%	44%	-	49%	44%	53% M	34%	48%	42%
A severe case of rotavirus can have more than 20 episodes of diarrhea or vomiting per day	162	24	27	64	36	12	55	107	162	0	10	152	99	63	72	90
	41%	38%	42%	46%	39%	33%	46%	39%	41%	-	35%	42%	47% M	35%	50% O	36%
Rotavirus often goes away without any medical treatment	137	17	18	44	46	10	35	102	137	0	5	132	87	50	62	75
	35%	28%	29%	32%	49% ABCE	29%	29%	37%	35%	-	18%	36%	41% M	28%	43% O	30%
Nearly all children will become infected with rotavirus by the time they enter kindergarten	121	15	14	51	29	10	44	77	121	0	11	110	81	40	58	63
	31%	25%	23%	38% B	31%	29%	37%	28%	31%	-	38%	30%	38% M	22%	40% O	25%
Rotavirus can be treated with an antibiotic	72	9	7	36	14	5	30	42	72	0	7	65	44	28	23	49
	18%	15%	11%	26% BD	15%	14%	25% G	15%	18%	-	24%	18%	21%	16%	16%	20%

Rotavirus Study (60-6056-01)

Q27. [SUMMARY - FALSE] Now, please indicate if you think the following statements about rotavirus or viral diarrhea are true or false.

Proportions/Means: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: Ever heard about rotavirus	394	61	62	136	93	42	121	273	394	0	29	365	212	182	144	250
Weighted	393	63*	64*	137	93*	37*	121	273	393	-**	29**	364	212	181	143	250
Rotavirus can be treated with an antibiotic	154	29	27	46	43	9	40	114	154	0	7	147	95	58	70	84
	39%	46% E	42%	34%	46% E	24%	33%	42%	39%	-	24%	40%	45% M	32%	49% O	33%
Rotavirus often goes away without any medical treatment	85	13	13	37	15	6	37	48	85	0	8	77	50	35	27	58
	22%	21%	21%	27%	16%	17%	31% G	18%	22%	-	28%	21%	24%	19%	19%	23%
Nearly all children will become infected with rotavirus by the time they enter kindergarten	82	10	12	25	31	3	20	62	82	0	4	78	40	42	21	61
	21%	16%	19%	18%	33% ACE	10%	17%	23%	21%	-	14%	22%	19%	23%	15%	25% N
Rotavirus is the most common cause of diarrhea in young children	43	8	6	17	8	3	17	26	43	0	2	41	26	17	16	27
	11%	13%	10%	13%	9%	10%	14%	10%	11%	-	7%	11%	12%	9%	11%	11%
A severe case of rotavirus can have more than 20 episodes of diarrhea or vomiting per day	14	3	0	5	6	0	5	9	14	0	1	13	10	4	5	9
	4%	5%	-	4%	6% B	-	4%	3%	4%	-	3%	4%	5%	2%	4%	4%
Rotavirus is contagious	10	2	5	3	0	0	4	6	10	0	1	9	5	5	4	6
	3%	3%	8% D	2%	-	-	3%	2%	3%	-	4%	3%	2%	3%	3%	2%
Rotavirus can be very serious and unpredictable	9	1	0	1	6	1	3	6	9	0	2	7	7	2	5	4
	2%	2%	-	1%	6% BC	2%	2%	2%	2%	-	7%	2%	3%	1%	4%	2%

Rotavirus Study (60-6056-01)

Q28. Besides your child's primary healthcare provider, from which two of the following sources do you obtain most of your information about health issues for your children?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Internet	475 58%	63 57%	72 50%	183 60%	116 58%	42 62%	150 57%	325 58%	235 60%	241 56%	15 52%	460 58%	146 61%	329 57%	181 61%	295 56%
Friend/ family member	470 57%	67 61%	83 58%	182 60%	100 50%	38 57%	160 61%	310 55%	198 50%	272 63%	18 62%	452 57%	117 49%	353 61%	172 58%	298 57%
Brochures or pamphlets	213 26%	22 20%	41 29%	73 24%	59 30%	18 27%	56 21%	157 28%	105 27%	108 25%	6 21%	207 26%	64 27%	149 26%	74 25%	139 27%
Television/ Radio	122 15%	12 11%	22 15%	43 14%	33 17%	11 17%	36 14%	86 15%	66 17%	55 13%	5 17%	117 15%	37 15%	85 15%	31 10%	91 17%
Magazines	116 14%	14 13%	28 19%	47 16%	18 9%	9 13%	40 15%	76 14%	76 19%	40 9%	5 17%	111 14%	47 20%	69 12%	34 11%	82 16%
Books	34 4%	2 2%	7 5%	15 5%	6 3%	3 5%	14 5%	20 4%	17 4%	17 4%	0 -	34 4%	11 5%	23 4%	17 6%	17 3%
Newspapers	34 4%	5 5%	5 4%	10 3%	10 5%	3 5%	11 4%	23 4%	19 5%	15 3%	2 7%	32 4%	10 4%	24 4%	12 4%	22 4%
CLSC/ Info-santé	26 3%	0 -	0 -	0 -	26 13% ABCE	0 -	10 4%	16 3%	7 2%	19 4% H	0 -	26 3%	7 3%	19 3%	13 4%	13 2%
Telehealth	22 3%	2 2%	5 4% D	15 5% DE	0 -	0 -	5 2%	17 3%	4 1%	18 4% H	0 -	22 3%	2 1%	20 3% L	13 4% O	9 2%
Nurse/ Public Health nurse	21 3%	8 7% CDE	8 6% CDE	3 1%	2 1%	0 -	10 4%	11 2%	11 3%	10 2%	1 4%	20 3%	7 3%	14 2%	9 3%	12 2%
Doctors	16 2%	1 1%	2 1%	5 2%	8 4%	0 -	3 1%	13 2%	13 3% I	3 1%	4 14%	12 2%	10 4% M	6 1%	6 2%	10 2%
B.C. Health Guide	8 1%	8 7% BCDE	0 -	0 -	0 -	0 -	1 0	7 1%	4 1%	4 1%	0 -	8 1%	4 2%	4 1%	6 2% O	2 0
Health unit/ public health unit	8 1%	2 2% C	4 3% CD	0 -	0 -	2 3% CD	2 1%	6 1%	4 1%	4 1%	0 -	8 1%	2 1%	6 1%	2 1%	6 1%
Pharmacists	7 1%	1 1%	0 -	2 1%	4 2%	0 -	4 2%	3 1%	4 1%	3 1%	0 -	7 1%	1 0	6 1%	3 1%	4 1%

Rotavirus Study (60-6056-01)

Q28. Besides your child's primary healthcare provider, from which two of the following sources do you obtain most of your information about health issues for your children?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Other	13 2%	2 2%	1 1%	6 2%	0 -	3 5% BD	5 2%	8 1%	7 2%	6 1%	0 -	13 2%	4 2%	9 2%	1 0	12 2% N
(DK/NS)	1 0	0 -	0 -	1 0	0 -	0 -	0 -	1 0	0 -	1 0	0 -	1 0	0 -	1 0	0 -	1 0

Rotavirus Study (60-6056-01)

Q29. Based on your two previous responses, can you name specific sources where you obtain your information about health issues for your children?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Friends/ family	307 37%	45 41%	58 41%	118 39%	62 31%	23 34%	97 37%	209 37%	140 36%	167 39%	13 44%	294 37%	88 37%	218 38%	118 40%	188 36%
Internet (unspecified)	167 20%	21 19%	22 15%	64 21%	53 27% BE	8 12%	49 19%	118 21%	84 21%	83 19%	4 14%	163 21%	55 23%	112 19%	68 23%	99 19%
Brochures/ pamphlets from doctor's office	88 11%	8 7%	25 17% AD	34 11%	16 8%	5 8%	24 9%	64 11%	40 10%	48 11%	1 3%	87 11%	23 10%	65 11%	26 9%	62 12%
CLSC Brochure/ CLSC/ info santé	72 9%	0 -	1 1%	4 1%	66 33% ABCE	1 1%	29 11%	43 8%	30 8%	42 10%	3 10%	69 9%	28 12%	44 8%	28 9%	44 8%
Family doctor	69 8%	9 8%	6 4%	28 9%	22 11% B	3 5%	26 10%	43 8%	48 12% I	21 5%	6 21%	63 8%	35 15% M	34 6%	20 7%	49 9%
Parent/ Parenting magazine	67 8%	10 9%	11 8%	23 8%	12 6%	10 16% CD	34 13% G	34 6%	50 13% I	17 4%	8 27%	59 7%	28 12% M	40 7%	23 8%	44 8%
Today's Parent	58 7%	6 6% D	10 7% D	35 12% D	0 -	6 9% D	15 6%	43 8%	41 10% I	17 4%	2 7%	56 7%	22 9%	36 6%	20 7%	38 7%
Telehealth	53 6%	1 1%	14 10% AD	32 11% AD	1 1%	4 6% AD	18 7%	35 6%	11 3%	42 10% H	1 3%	52 7%	9 4%	44 8% L	26 9% O	27 5%
Google	47 6%	7 7%	6 4%	25 8% D	5 3%	3 5%	16 6%	31 6%	27 7%	20 5%	2 7%	45 6%	11 5%	36 6%	17 6%	30 6%
Brochures/ information from the pharmacy	41 5%	4 4%	4 3%	13 4%	14 7%	5 8%	10 4%	31 5%	20 5%	21 5%	1 3%	40 5%	8 3%	33 6%	15 5%	26 5%
Nurse	40 5%	11 10% CD	15 11% CD	5 2%	6 3%	3 4%	16 6%	24 4%	19 5%	21 5%	2 7%	38 5%	9 4%	31 5%	14 5%	26 5%
The news (unspecified)	40 5%	5 5%	8 6%	15 5%	7 4%	4 6%	11 4%	29 5%	15 4%	25 6%	2 7%	38 5%	9 4%	31 5%	9 3%	31 6%

Rotavirus Study (60-6056-01)

Q29. Based on your two previous responses, can you name specific sources where you obtain your information about health issues for your children?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O
Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
webMD.com	36 4%	9 8% D	7 5% D	16 5% D	0 - D	3 5% D	9 3%	27 5%	19 5%	17 4%	1 4%	35 4%	7 3%	29 5%	16 5%	20 4%
Babycenter.com	33 4%	5 5% D	7 5% D	19 6% D	0 - D	2 3% D	13 5%	20 4%	19 5%	14 3%	1 4%	32 4%	13 5%	20 3%	13 4%	20 4%
Differents medical books on baby's,childrens	33 4%	3 3%	6 4%	13 4%	7 4%	3 5%	14 5%	19 3%	18 5%	15 3%	2 7%	31 4%	10 4%	23 4%	17 6%	16 3%
The health unit/public health/ clinic/ center	32 4%	7 7% CD	12 9% CD	7 2%	2 1%	3 5% D	7 3%	25 4%	16 4%	16 4%	2 7%	30 4%	8 3%	24 4%	12 4%	20 4%
Television (unspecified)	30 4%	1 1%	3 2%	11 4%	10 5%	4 6% A	11 4%	19 3%	18 5%	12 3%	0 -	30 4%	13 5%	16 3%	7 2%	23 4%
Health Canada (web site,brochure,etc...)	26 3%	4 4%	2 1%	14 5%	3 2%	3 4%	11 4%	15 3%	11 3%	15 3%	1 3%	25 3%	5 2%	21 4%	9 3%	17 3%
News papers	23 3%	6 6% C	2 1%	5 2%	7 4%	3 4%	7 3%	16 3%	12 3%	11 3%	0 -	23 3%	7 3%	16 3%	7 2%	16 3%
What to expect book (give by the hospital)	22 3%	2 2%	2 1%	4 1%	12 6% BC	2 3%	9 3%	13 2%	8 2%	14 3%	0 -	22 3%	7 3%	15 3%	10 3%	12 2%
B.C Health Guide	16 2%	15 14% BCDE	0 -	1 0	0 -	0 -	6 2%	10 2%	7 2%	9 2%	0 -	16 2%	5 2%	11 2%	9 3%	7 1%
Radio	10 1%	2 2%	2 1%	5 2%	1 1%	0 -	1 0	9 2%	3 1%	7 2%	0 -	10 1%	5 2%	5 1%	3 1%	7 1%
Mayo clinic.com	9 1%	2 2%	2 1%	3 1%	0 -	2 3% D	2 1%	7 1%	4 1%	5 1%	0 -	9 1%	1 0	8 1%	3 1%	6 1%
At the hospital	8 1%	1 1%	2 1%	3 1%	2 1%	0 -	3 1%	5 1%	1 0	7 2% H	0 -	8 1%	3 1%	5 1%	5 2%	3 1%
Mamanpourlaviewwebsite.com	8 1%	0 -	0 -	0 -	8 4% ABC	0 -	5 2%	3 1%	5 1%	3 1%	0 -	8 1%	3 1%	5 1%	3 1%	5 1%
Doctissimo	8 1%	0 -	0 -	1 0	6 3% BC	1 1%	1 0	7 1%	3 1%	5 1%	0 -	8 1%	4 2%	4 1%	4 1%	4 1%
Pampers	7 1%	0 -	1 1%	3 1%	2 1%	1 1%	3 1%	4 1%	2 0	5 1%	0 -	7 1%	1 0	6 1%	4 1%	3 1%

Rotavirus Study (60-6056-01)

Q29. Based on your two previous responses, can you name specific sources where you obtain your information about health issues for your children?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Babyzone.com	6 1%	3 3% CD	2 1%	1 0	0 -	0 -	1 0	5 1%	3 1%	3 1%	0 -	6 1%	3 1%	3 1%	3 1%	3 1%
Word of mouth	6 1%	0 -	0 -	4 1%	0 -	2 3% BD	2 1%	4 1%	1 0	5 1%	0 -	6 1%	1 0	5 1%	2 1%	4 1%
Other	85 10%	14 13%	13 9%	31 10%	15 8%	11 17% D	31 12%	54 10%	45 12%	40 9%	3 11%	82 10%	23 9%	63 11%	29 10%	56 11%
None	11 1%	1 1%	3 2%	3 1%	3 2%	1 1%	1 0	10 2%	6 2%	5 1%	2 7%	9 1%	3 1%	8 1%	3 1%	8 2%
Dk/ ns	22 3%	1 1%	6 4% D	11 4% D	1 1%	3 4% D	7 3%	15 3%	9 2%	13 3%	0 -	22 3%	4 2%	18 3%	4 1%	18 3%

Rotavirus Study (60-6056-01)

Q30. How would you describe the area where you live -- is it located in a city, in a suburb outside of a city, in a small town, or in a country or farm area?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
Base: All respondents Weighted		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
In a city	438 53%	47 43%	91 64% ADE	179 59% AE	100 50% E	20 30%	149 57%	289 52%	205 52%	233 54%	14 49%	423 53%	128 53%	309 53%	148 50%	289 55%
In a suburb outside of a city	160 19%	32 29% BE	9 6%	62 20% B	47 24% B	10 16% B	46 18%	114 20%	83 21%	77 18%	4 13%	156 20%	54 23%	106 18%	62 21%	98 19%
In a small town	173 21%	26 23%	27 19%	54 18%	40 20%	26 39% ABCD	51 19%	123 22%	78 20%	95 22%	9 31%	164 21%	45 19%	128 22%	67 22%	106 20%
In a country or farm area	44 5%	5 5%	12 9% C	5 2%	11 6% C	10 16% ACD	15 6%	29 5%	23 6%	21 5%	2 7%	42 5%	11 4%	33 6%	16 5%	28 5%
Small village/ small community	5 1%	0 -	2 1%	3 1%	0 -	0 -	1 0	4 1%	3 1%	2 0	0 -	5 1%	2 1%	3 1%	4 1% O	1 0
Other	2 0	0 -	1 1%	0 -	1 1%	0 -	0 -	2 0	1 0	1 0	0 -	2 0	0 -	2 0	1 0	1 0

Rotavirus Study (60-6056-01)

Q14/16. Has your child ever experienced five or more episodes of profuse, watery diarrhea in one day?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents Weighted	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Yes	299	45	46	100	80	27	66	233	143	155	10	289	104	194	299	0
	36%	41%	32%	33%	40%	40%	25%	42% F	36%	36%	34%	36%	43% M	33%	100% O	-
No	523	65	96	203	119	40	196	327	250	273	19	504	136	387	0	523
	64%	59%	68%	67%	60%	60%	75% G	58%	64%	64%	66%	64%	57% L	67%	-	100% N

Rotavirus Study (60-6056-01)

Q24/Q25. Has your child ever been diagnosed by a doctor with viral diarrhea or rotavirus?

Proportions/Mean: Columns Tested (5% risk level) - A/B/C/D/E - F/G - H/I - J/K - L/M - N/O

Overlap formulae used. * small base; ** very small base (under 30) ineligible for sig testing

		Region					Number of Child in Household		Heard of Rotavirus		Received Rotavirus Vaccine		Heard something about Rotavirus		Child/children experienced rotavirus	
	Total	BC	PR	Ont	Qc	Atl	1	2+	Yes	No	Yes	No	Yes	No	Yes	No
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
Base: All respondents	822	107	139	300	199	77	262	560	394	428	29	793	240	582	299	523
Weighted	822	110	143	303	200	67*	262	560	393	429	29**	793	240	582	299	523
Yes	85 10%	10 9%	11 8%	25 8%	34 17% BCE	4 6%	17 7%	68 12% F	55 14% I	30 7%	3 10%	82 10%	48 20% M	37 6%	70 24% O	15 3%
No	737 90%	100 91%	131 92% D	278 92% D	166 83%	63 94% D	245 93% G	492 88%	338 86%	399 93% H	26 90%	711 90%	192 80%	545 94% L	228 76%	508 97% N