

Q11. Which of the following have you experienced in the last year?

		Age			Gender		Split Sample Variable		Current Situation		Insurance/Benefits				Familiarity Chiropractors		Choice		Overall impression	
	Total	18-34	35-54	55+	Male	Female	Chiropractor	Neuromusculoskeletal Expert	Currently seeing chiropractor	Discontinued seeing chiropractor	Employer Benefits (per health practitioner)	Employer Benefits (spending account)	Private Insurance	None	Familiar	Unfamiliar	Chiropractor	Neuromusculoskeletal Expert	Favourable	Unfavourable
		A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S
Base: All respondents	3017	842	1218	957	1491	1526	1509	1508	223	2794	1221	332	380	1189	1873	1144	1490	1527	2007	728
Weighted	3017	841	1116	1060	1448	1569	1514	1503	227	2790	1183	313	375	1240	1867	1150	1474	1543	1991	737
Back pain and stiffness	1596	427	619	550	744	852	803	793	168	1428	704	155	182	611	1101	495	797	799	1109	377
	53%	51%	55%	52%	51%	54%	53%	53%	74%	51%	60%	49%	48%	49%	59%	43%	54%	52%	56%	51%
Headaches/migraine	1316	483	556	278	478	839	650	666	111	1206	565	137	126	541	843	473	675	641	914	302
	44%	57%	50%	26%	33%	53%	43%	44%	49%	43%	48%	44%	34%	44%	45%	41%	46%	42%	46%	41%
		BC	C			D					L	L		L	O		Q		S	
Neck pain	1069	321	418	329	453	615	535	534	128	940	464	111	121	410	766	302	549	520	768	232
	35%	38%	37%	31%	31%	39%	35%	36%	57%	34%	39%	35%	32%	33%	41%	26%	37%	34%	39%	31%
Joint pain		C	C			D			I		LM				O				S	
	1050	199	401	450	504	546	526	524	91	959	452	99	140	399	751	300	479	571	706	268
	35%	24%	36%	43%	35%	35%	35%	35%	40%	34%	38%	32%	37%	32%	40%	26%	33%	37%	36%	36%
Knee pain or injury		A	AB				KM								O					
	920	194	343	383	416	504	465	455	87	833	375	108	111	360	638	282	439	481	623	226
Shoulder pain or injury		31%	23%	31%	36%	29%	31%	30%	38%	30%	32%	35%	30%	29%	34%	25%	30%	31%	31%	31%
		A	AB				I								O					
	870	195	360	315	407	463	438	431	111	759	396	82	95	323	638	232	432	437	602	211
Muscle spasms or tension		29%	23%	32%	30%	28%	29%	29%	49%	27%	34%	26%	25%	26%	34%	20%	29%	28%	30%	29%
		A	A				I				KLM				O					
	824	239	336	248	354	470	408	416	91	733	364	90	98	307	595	229	396	428	563	198
Arthritis		27%	28%	30%	23%	24%	30%	27%	40%	26%	31%	29%	26%	25%	32%	20%	27%	28%	28%	27%
		C	C			D			I		M				O					
	672	47	227	398	317	355	327	346	56	616	240	57	111	278	472	200	298	374	457	162
Hip pain or injury		22%	6%	20%	38%	22%	22%	23%	25%	22%	20%	18%	30%	22%	25%	17%	20%	24%	23%	22%
		A	AB												O			P		
	515	99	196	219	193	322	267	248	64	450	206	43	69	220	380	134	261	254	345	128
Muscle weakness		17%	12%	18%	21%	13%	18%	17%	28%	16%	17%	14%	19%	18%	20%	12%	18%	17%	17%	17%
		A	A			D			I						O					
	454	128	169	158	209	246	231	223	50	405	174	44	47	203	335	120	203	251	303	122
Pinched nerves		15%	15%	15%	14%	16%	15%	15%	22%	15%	15%	14%	13%	16%	18%	10%	14%	16%	15%	17%
							I								O					
	404	93	160	151	185	219	195	209	54	350	166	39	49	165	308	96	208	196	287	98
Tendonitis		13%	11%	14%	14%	13%	13%	14%	24%	13%	14%	13%	13%	13%	17%	8%	14%	13%	14%	13%
		A					I								O					
	320	68	134	119	131	189	150	170	43	277	140	42	31	119	225	95	153	168	218	77
Sport-related injury		11%	8%	12%	11%	9%	10%	11%	19%	10%	12%	14%	8%	10%	12%	8%	10%	11%	11%	10%
		A	A			D			I						O					
	273	104	114	55	181	93	123	150	33	240	153	40	28	72	197	76	126	147	165	86
Spinal problem		9%	12%	10%	5%	6%	8%	10%	15%	9%	13%	13%	7%	6%	11%	7%	9%	10%	8%	12%
		C	C		E				I		LM	LM			O				R	
	236	30	87	118	115	120	118	118	43	193	97	19	34	96	181	55	103	133	149	68
Disc disease or herniation		8%	4%	8%	11%	8%	8%	8%	19%	7%	8%	6%	9%	8%	10%	5%	7%	9%	8%	9%
		A	AB				I								O					
	159	15	57	88	84	76	78	82	22	138	71	19	18	59	121	38	60	100	94	54
Scoliosis		5%	2%	5%	8%	6%	5%	5%	10%	5%	6%	6%	5%	5%	7%	3%	4%	7%	5%	7%
		A	AB				I								O			P		R
	79	24	25	30	27	52	41	38	16	63	37	7	7	30	58	21	33	46	52	20
Whiplash		3%	3%	2%	3%	2%	3%	3%	7%	2%	3%	2%	2%	2%	3%	2%	2%	3%	3%	3%
						D			I						O					
	68	34	23	11	25	43	38	30	14	54	28	8	11	27	48	21	40	29	46	13
Other		2%	4%	2%	1%	2%	3%	2%	6%	2%	2%	3%	3%	2%	3%	2%	3%	2%	2%	2%
		BC							I											
	307	68	111	128	158	150	147	160	11	296	99	27	46	143	154	153	128	180	174	66
Summary		10%	8%	10%	12%	11%	10%	11%	5%	11%	8%	9%	12%	12%	8%	13%	9%	12%	9%	9%
		A								H			J	J		N		P		
Only 1 symptom	888	260	329	299	482	406	444	444	22	866	273	91	125	420	393	495	447	441	540	205
	29%	31%	30%	28%	33%	26%	29%	30%	10%	31%	23%	29%	33%	34%	21%	43%	30%	29%	27%	28%
2-3 symptoms		E							H			J	J		N					
	846	269	277	300	405	441	429	417	54	792	338	88	94	348	542	304	405	441	561	224
	28%	32%	25%	28%	28%	28%	28%	28%	24%	28%	23%	28%	25%	28%	29%	27%	28%	29%	28%	30%
4-5 symptoms		B																		
	579	162	221	195	272	306	301	278	58	521	264	76	75	189	386	193	286	293	394	150
	19%	19%	20%	18%	19%	20%	20%	19%	26%	19%	22%	24%	20%	15%	21%	17%	19%	19%	20%	20%
6+ symptoms		I									M	M			O					
	704	150	289	265	289	415	340	364	93	611	308	58	81	284	546	158	337	367	496	158
	23%	18%	26%	25%	20%	27%	22%	24%	41%	22%	26%	18%	22%	23%	29%	14%	23%	24%	25%	21%
		A	A			D			I		K				O					

Proportions/Mean: Columns Tested (5% risk level) - A/B/C - D/E - F/G - H/I - J/K/L/M - N/O - P/Q - R/S Overlap formulae used.